

Antioxidants, Radical Scavenging, and Their Impact on Oxidative Stress

Franklyn Okechukwu Ohiagu¹, Paul Chidoka Chikezie^{2,3,*}, Tochukwu Oluwatosin Maduka⁴, Chinwendu Mauren Chikezie⁵, Ogueri Nwaiwu⁶, Keshav Raj Paudel⁷

¹Department of Biochemistry, Federal University of Technology, Owerri, NIGERIA.

²School of Sciences, Coventry University, Coventry, England, UNITED KINGDOM.

³Department of Biochemistry, Imo State University, Owerri, NIGERIA.

⁴Graduate School of Science and Engineering, Saitama University, Saitama City, JAPAN.

⁵Department of Biomedical Sciences, University of Hull, Kingston upon Hull, UNITED KINGDOM.

⁶School of Chemical Engineering, University of Birmingham, Edgbaston, UNITED KINGDOM.

⁷School of Life Sciences, University of Technology, Sydney, AUSTRALIA.

ABSTRACT

Antioxidants impede free radical chain reactions and thereby retard and prevent oxidative cellular damage. The present review provides an overview of the basic mechanisms by which antioxidants scavenge free radicals and describes the basic methods for measurement of antioxidant activities against free radicals as well as the underlying chemistry and pathologies linked with oxidative stress. This review was sourced online from scientific search engines using appropriate keywords. For appropriate measurement of antioxidant capacity, with respect to reaction mechanisms involved in the free radical reduction process, exclusive application of HAT, SET methods, and/or a combination of both HAT/SET methods should be considered. The human body is equipped with an antioxidant defence system to mitigate or prevent free radical-induced cellular damage through free radical scavenging mechanisms such as inhibition of enzyme activity involved in free radical generation, activation of intracellular enzymatic antioxidant activity, metal ion chelation, as well as inhibition of protein modification, DNA damage, and lipid peroxidation. Impaired antioxidant activity is associated with chronic diseases such as cardiovascular diseases, diabetes mellitus, inflammatory disease, cancer, cataracts, Alzheimer's disease, autism, and ageing. Overexpression of enzymatic antioxidants, as demonstrated in mutant mice models, could serve as a novel alleviative measure against the development of pathological conditions associated with oxidative stress.

Keywords: Antioxidants, Enzymatic antioxidants, Free radical, Lipid peroxidation, Oxidative stress.

Correspondence:

Dr. Paul Chidoka Chikezie

School of Sciences, Coventry University,
Coventry, England, UNITED KINGDOM.

Department of Biochemistry, Imo State
University, Owerri, NIGERIA.

Email: ae2389@coventry.ac.uk

Received: 27-12-2024;

Revised: 30-12-2024;

Accepted: 07-01-2025.

INTRODUCTION

Antioxidants impede free radical chain reactions and thereby retard and prevent oxidative cellular damage. Although antioxidant defences differ in diverse species, the antioxidant defence mechanism is universal. Antioxidants are enzymatic or non-enzymatic entities present in extracellular and intracellular environments.¹ Antioxidants are seen occurring naturally in plants, animals, and microorganisms or are synthesised using chemical methods. Higher plant constituents are rich sources of natural antioxidants, such as the polyphenols and tocopherols, which occur in large quantities in fruits, herbs, vegetables, spices, cereals, oils, grains, and seeds. Marine organisms, such as fish/

shellfish, algae, and marine bacteria, were also reported as sources of antioxidants.^{2,3} Furthermore, food and agricultural industrial byproducts, such as skins and shells of nuts, as well as cereals, fish viscera extracts, citrus peels, etc., have been considered as rich sources of antioxidants.³⁻⁶ It is worthwhile to note that regular intake of dietary antioxidants is essential to sustain adequate antioxidant status.^{7,8}

Free radicals are referred to as atoms, molecules, or ions that possess unpaired electrons and react very actively with other molecules. Biologically, free radicals arise from oxygen, sulphur, and nitrogen molecules and form parts of groups of molecules known as ROS, RSS, and RNS.⁹

High cellular levels of ROS cause harmful effects on various biological molecules such as protein, lipid, DNA, and RNA. Evidence has shown that ROS attack on macromolecules is one of several attributes of oxidative stress. Oxidative stress contributes to the development of CVD, DM, inflammatory disease, cancer,



DOI: 10.5530/fra.2024.2.7

Copyright Information :

Copyright Author (s) 2024 Distributed under
Creative Commons CC-BY 4.0

Publishing Partner : Manuscript Technomedia. [www.mstechnomedia.com]

cataracts, AD, autism, and aging. The ROS levels usually increase in conditions of environmental stress and cell dysfunction, eliciting considerable damage to body cells.¹⁰⁻¹²

The human body consists of an antioxidant defence system for the mitigation or prevention of free radical-induced cellular damage through free radical scavenging mechanisms, such as inhibition of enzyme activity involved in free radical generation, activation of intracellular enzymatic antioxidant activity, metal ion chelation, as well as inhibition of protein modification, DNA damage, and lipid peroxidation. Antioxidants neutralise free radicals by donating or accepting electron(s) to obliterate unpaired electrons of radicals. Antioxidants neutralise reactive radicals such that the radicals are less active and deleterious with longer half-lives compared with the neutralised radicals. In addition, free radicals are further neutralised by the actions of other antioxidants or through other mechanisms that annihilate their deleterious activities.⁸

Antioxidant activity assays have undergone remarkable advancement within the last few decades. Chemical assays that are made up of high sensitivity as well as automated detection technologies are used to evaluate antioxidant activity through mechanisms such as free radical scavenging activity, HAT, SET, reducing power, etc.,^{13,14} A good knowledge of the basic mechanisms, usefulness, and limitations of the various assays is essential for the selection of appropriate method(s) required for measurement of antioxidant capacity of desired application.¹⁴

The mode of actions of radicals and mechanisms of antioxidant activities, in addition to the connecting cellular events leading to oxidative stress, relate to the pathogenesis and pathophysiology of human disorders as well as the ageing process. And in that regard, measurements of cellular redox status and perturbation levels are of immense application in prognosis/diagnosis and mitigation strategies against human pathological conditions. Therefore, the present review provides an overview of the basic mechanisms by which antioxidants scavenge free radicals and describes the basic methods for measurement of antioxidant activities against free radicals as well as the underlying chemistry. The pathophysiology of oxidative stress-induced states, as well as establishing novel scientific approaches to prevention and mitigation of oxidative stress-induced pathological conditions, were also highlighted.

Evidence Acquisition

This review was sourced online from scientific search engines, namely, ResearchGate, Google Scholar, Scopus, PubMed, Medline and SpringerLink, using keywords, namely, 'antioxidants', 'enzymatic antioxidants', 'electron transfer', 'free radical', 'hydrogen atom transfer', 'oxygen radical', 'reactive oxygen species', 'reactive sulphur species', 'reactive nitrogen species', 'metal ion chelating', 'lipid peroxidation', 'DNA damage', 'aging', 'Alzheimer's disease', 'Parkinson's disease', 'cancer', 'diabetes mellitus', 'cardiovascular diseases', 'sickle cell disease', 'oxidative stress and pathology'. A

total number of 283 references published online between 1959 and 2022 were used as information sources and cited in this review.

BASES FOR ANTIOXIDANTS CATEGORIES

Antioxidants are categorised based on their activity, solubility, size, mode of action, and occurrence. Based on activity, antioxidants can be categorised as 'enzymatic' and 'non-enzymatic' entities. The enzymatic antioxidants can convert deleterious oxidative products to H₂O₂ and finally to water, as well as possessing the capacity to break down and eliminate free radicals using cofactors such as Fe, Zn, Cu, Se, and Mn. The non-enzymatic antioxidants are ascorbic acid, α -tocopherol, carotenoids, glutathione, as well as plant polyphenols. These antioxidants act to hinder and interfere with free radical chain reactions.^{1,14,15}

Based on solubility, there are 'water-soluble' and 'lipid-soluble' antioxidants. For example, ascorbic acid and α -tocopherol are water-soluble and lipid-soluble antioxidants, respectively.¹⁵ Based on size, antioxidants are classified as 'small-molecule' and 'large-molecule.' The LMWAs include the carotenoids, ascorbic acid, α -tocopherol, glutathione, etc., The HMWA are enzymes such as CAT, SOD, and GPx, as well as sacrificial proteins (albumin), which absorb ROS and inhibit deleterious actions of radicals against important macromolecules.^{1,14}




Based on mode of action, antioxidants are categorised into 6 groups, namely: (1) antioxidants that break chain reactions by reacting with RO₂[•] that are composed of weak O-H or N-H bonds as characteristics of the phenols, aromatic amines, naphthols, aminophenols, and hydroquinones. (2) Antioxidants that possess the capacity to break free radical chain reactions by reacting with alkyl radicals. And these include the quinones, iminoquinones, and nitrones. (3) There are antioxidants that terminate cyclic free radical chain reactions of nitroxyl radicals, aromatic amines, and variable valence metal compounds. (4) Another form of antioxidants that disintegrates 'OOH are the phosphides, thiophosphates, and sulfides. (5) Antioxidants that deactivate hydroxyl acids, diamines, and bifunctional compounds. (6) Finally, the phenol sulphide antioxidants, whose phenolic group reacts with RO₂[•] of sulphide group and 'OOH in a synergistic manner.^{1,15}

Based on occurrence, antioxidants are classified as 'natural' or 'synthetic' products. The natural antioxidants are chain-breaking antioxidants that undergo reactions with radicals and convert them to more stable adducts. The natural antioxidants are phenolic in structure and include the vitamins, phytochemicals, and minerals (which are enzymatic antioxidants cofactors such like Mn, Fe, Cu, Zn, and Se. The synthetic antioxidants are also phenolic compound derivatives that trap free radicals and terminate free radical chain reactions. The synthetic antioxidants include BHT, metal chelating agent-EDTA, NDGA, BHATBHQ, and PG.^{16,17}



SOURCES OF FREE RADICALS

The ROS are mainly generated in the mitochondria, through the electron transport chain events associated with aerobic respiration.¹⁸ The ROS include $O_2^{\cdot-}$, 1O_2 , $\cdot OH$, $RO\cdot$, $RO_2^{\cdot-}$, H_2O_2 , $LOOH$.^{1,19} Approximately 1-3% of electrons react prematurely with oxygen during the electron transport chain to generate $O_2^{\cdot-}$.²⁰

The first mechanism involved in ROS generation is the formation of $O_2^{\cdot-}$ from oxygen and NADPH by the action of NADPH oxidase (Eqn 1). The $O_2^{\cdot-}$ is further acted upon by SOD to generate H_2O_2 (Eqn 2). The MPO in the neutrophil cytoplasmic granules converts H_2O_2 in the presence of Cl^- to $HOCl$ (Eqn 3).²¹



The $O_2^{\cdot-}$ and H_2O_2 are also involved in other ROS generation suchlike $\cdot OH$ through respiratory burst by Fenton (Eqn 4) and/or Haber-Weiss reactions (Eqn 5).²²



The RNS suchlike $NO\cdot$ is generated by eNOS (Eqn 6). The reaction between $NO\cdot$ and $O_2^{\cdot-}$ leads to the production of a very strong oxidant- $ONOO\cdot$.²³



ROS and RNS are generated endogenously through inflammation mechanisms and immune cell stimulation, ageing, extreme exercise, mental stress, ischaemia, infectious diseases, etc. Furthermore, ROS and RNS emanate exogenously through smoking, the presence of heavy metals, and exposure to radiation, as well as the intake of alcohol, some drugs such as cyclosporine and tacrolimus, and certain organic solvents such as benzene, polluted air inhalation, polluted water consumption, etc., These compounds, often referred to as pro-oxidants, are converted to ROS in the body system.²⁴ The ROS are also generated via lipo-oxygenation, lipid peroxidation, and cyclo-oxygenation.¹⁴

Mechanisms of Action of Antioxidants Against Free Radicals

Antioxidants scavenge free radicals and impede their actions through inhibition of enzymes involved in free radical production, activation of intracellular enzymatic antioxidants, metal ion chelation, and inhibition of protein modification, DNA damage, and lipid peroxidation.

Inhibition of Enzymes Involved in Free Radical Generation



A reasonable number of enzymes are involved in various free radical generation. The plasma membrane-linked enzymes NADPH oxidases induce $O_2^{\cdot-}$ formation by transferring one electron from the cytosolic donor NADPH to a molecule of molecular oxygen.²⁵ Xanthine oxidase enhances $O_2^{\cdot-}$ and H_2O_2 production through the oxidation of hypoxanthine and xanthine to uric acid.²⁶ The $O_2^{\cdot-}$ is also produced by enzymes such as cyclooxygenases, monooxygenases, and NADH oxidase. Excessive levels of $O_2^{\cdot-}$ are deleterious. However, it offers protection to the body against invading microorganisms during oxygen-dependent pathogen obliteration mechanisms.²⁷

Many natural antioxidants are potential inhibitors of enzymes that promote $O_2^{\cdot-}$ generation and other ROS. For instance, polyphenols and α -tocopherol are known inhibitors of NADPH oxidase activity.²⁸ In addition; the polyphenolic compound coumarin elicits an inhibitory effect on xanthine oxidase.²⁹


Intracellular Enzymatic Antioxidants

The ROS, namely $O_2^{\cdot-}$, $\cdot OH$, H_2O_2 , etc., are irreversibly generated during metabolic events. Certain intracellular enzymes, such as SOD, GST, CAT, GR, GP_x , TrxR, biliverdin reductase, and heme oxygenase, extensively mitigate the levels of ROS.^{8,30}


The SOD enzymes (which are metal-containing proteins) catalyse the dismutation of two molecules of $O_2^{\cdot-}$ to O_2 and the less reactive species, H_2O_2 (Eqns 7 and 8; Figure 1).^{8,31} The survival rate of rats subjected to 100% oxygenation increased following intravenous injection of liposomes containing SOD and CAT before and during the exposure to oxygen.³² According to the study by Li *et al.*³³ lack of expression of Mn-SOD in Mn-SOD knockout mice was accompanied by dilated cardiomyopathy as well as neonatal lethality. Van Loon *et al.*³⁴ also reported that Mn-SOD plays an important role in the natural protection of cells against oxygen toxicity in their study using yeast mutants.




The H_2O_2 can be further neutralized to form water via the reactions catalyzed by CAT or GP_x . Thus, CAT and/or GP_x catalyze the eradication of H_2O_2 , and thereby prevent $\cdot OH$ generation through the Fenton Reaction.



The GP_x uses hydrogen from two molecules of GSH to nullify H_2O_2 to two water molecules, forming GSSG in the process (Eqns 9 and 10; Figure 1).^{8,31,35}



The GR further catalyzes the reduction of the GSSG back to GSH, thereby completing the cycle (Eqn 11; Figure 1).^{31,36}



Metal Ion Chelating

Transition metals, such as copper and iron, exacerbate oxidative stress. Fe^{2+} and Cu^+ ions undergo a reaction with H_2O_2 generated from dismutation of $\text{O}_2\cdot^-$ by SOD to form a very reactive free radical, $\cdot\text{OH}$ (Eqn 12). In comparison with iron, copper produces more singlet oxygen than $\cdot\text{OH}$ when reacted with H_2O_2 . During this process, Fe^{2+} and Cu^+ ions undergo oxidation to Fe^{3+} and Cu^{2+} ions, respectively. Furthermore, the Fe^{3+} and Cu^{2+} ions formed can undergo reduction back to Fe^{2+} and Cu^+ ions in the presence of reducing agents such as AscH. Thus, Fe^{2+} and Cu^+ ions can again react with another H_2O_2 molecule to produce more $\cdot\text{OH}$ (Eqn 13).^{8,15}

The $\cdot\text{OH}$ generated can directly react vigorously with proteins and other macromolecules to form carbonyls, namely, aldehydes and ketones, leading to cross-linking and lipid peroxidation. Thus, the formation and action of $\cdot\text{OH}$ and ROS is impeded through chelation of metal ions, especially iron and copper.⁸



Metal chelating is often regarded as the most acknowledged and common antioxidant action/mechanism. Antioxidants bind to metal ions through their functional groups, as observed in Fe binding to an antioxidant such as taxifolin. The chelating sites for the Fe^{2+} ion to taxifolin is the 4-oxo and 5-OH groups that are found between the heterocyclic and the A rings, the catechol moiety of the B ring, and the 3-OH groups and 4-oxo group in the heterocyclic ring (Figure 2A). Thus, taxifolin can chelate more than one Fe^{2+} ion through its functional hydroxyl and carbonyl groups.³⁷ Other functional groups through which antioxidants chelate metal ions, such as Fe^{2+} and Cu^{2+} ions, include -OH and - OCH_3 groups as observed in curcumin and kaempferol.³⁸ This metal-chelating capability of antioxidants was reported in the study carried out by Kazazica *et al.*³⁹ in which kaempferol was bonded to Fe^{2+} and Cu^{2+} ions through -OH and - OCH_3 groups (Figure 2D). Another antioxidant, usnic, was reported to chelate Fe^{2+} ions through its -OH and -COOH groups bonded to the phenolic ring (Figure 2B).⁴⁰ In another study, the antioxidant capability of resveratrol was linked to its metal-chelating property, whereby Fe^{2+} ions are chelated through their -OH groups at the meta positions (Figure 2C).⁴¹ *In vitro* studies on metal ion chelating efficacy by antioxidants showed that quercetin, (+) catechin, and rutin efficiently chelated Fe^{2+} ions.^{42,43}

Other functional groups through which antioxidants chelate metal ions, such as Fe^{2+} and Cu^{2+} ions, include -OH and - OCH_3

groups as observed in curcumin and kaempferol (Figure 2D).^{38,39} Another antioxidant, usnic, has been reported to chelate Fe^{2+} ions through its -OH and -COOH groups bonded to the phenolic ring (Figure 2B).⁴⁰ The antioxidant capacity of resveratrol has been linked to its metal-chelating capacity, as they have been reported to chelate Fe^{2+} ions via their -OH groups at meta positions (Figure 2C).⁴¹

Generally, compounds made up of two or more -COOH, -C=O, -OH, - OCH_3 , -SH, - PO_3H_2 , -O-, - NR_2 , and -S- functional groups in the suitable structural conformation can effectively chelate Fe^{2+} ions.⁴⁴⁻⁴⁷ However, the interaction between antioxidants and metal ions can disrupt the biological actions and properties of antioxidants.^{48,49} Metal chelation is also an important aspect of the antioxidant mechanism that mitigates the concentration of metals involved in lipid peroxidation catalytic activities.⁵⁰

Inhibition of Protein Modification

Free radicals initiate the modification of proteins through amino acid nitration or chlorination. The RNS, namely, ONOO^- , is a known powerful oxidant and nitrating agent.⁵² The ONOO^- has the capacity to initiate considerable damage to cellular proteins and DNA. Both ONOO^- and its protonated form, ONOOH , modify proteins via one- or two-electron oxidation processes.⁵³ The ONOO^- undergoes reaction with carbon IV oxide yielding ONOOCO_2^- . The ONOOCO_2^- decomposes, leading to $\text{CO}_3\cdot^-$ as well as $\cdot\text{NO}_2$ generation. The $\cdot\text{NO}_2$ is a RNS that has the capacity to nitrate Tyr to nitrotyrosine. Thus, these radicals are known to elicit ONOO^- -related cellular damage (Figure 3). Furthermore, ONOO^- can directly undergo reactions with electron-rich groups such as iron-sulphur centres, zinc-thiolates, sulfhydryls, as well as the sulfhydryl active site that occurs in tyrosine phosphatases.⁵²

An alternative pathway of protein nitration or chlorination involves heme protein MPO-generated HOCl, which undergoes reaction with NO_2^- to generate NO_2Cl . The NO_2Cl may further disintegrate spontaneously to yield NO_2^- and Cl^\bullet . The NO_2^- and Cl^\bullet are probably responsible for the nitrating and chlorinating capacities of NO_2Cl .⁵⁴ Chlorotyrosine and nitrotyrosine are biomarkers for disorders caused by RNS.⁵⁵

These modifications usually lead to changes in the function and structure of proteins, especially inhibition of enzyme activities. Tissue proteins made up of nitrotyrosine residues have been reported to be present in various pathologic conditions such as hypertension, atherosclerosis, and DM and are associated with increased ONOO^- generation.⁵⁶

Non-enzymatic antioxidants and enzymatic antioxidants suppress protein modification elicited by ONOO^- and HOCl. Elimination of H_2O_2 by antioxidants, such as the enzymatic antioxidants, CAT, will inhibit HOCl generation. Furthermore, antioxidants such as polyphenols, curcumin, and the enzymatic antioxidant SOD suppress ONOO^- formation through $\text{O}_2\cdot^-$ scavenging (Figure

3).⁵⁷ Both *in vivo* and *in vitro* studies on the antioxidant efficacy of S-allylcysteine revealed the capability of the garlic-derived compound to scavenge HOCl, O₂^{•-}, H₂O₂, [•]OH and ONOO⁻.⁵⁸ Furthermore, Fernandes *et al.*⁵⁹ reported that O₂^{•-}, [•]OH, NO[•], and ONOO⁻ are readily scavenged by sulindac and sulindac sulfone, whereas sulindac sulphide scavenged HOCl, O₂^{•-}, [•]OH, NO[•], and ONOO⁻.

Inhibition of DNA Damage

The [•]OH and ONOO⁻ react with plasmid DNA and nick or cleave one DNA strand, resulting in the unwinding of the supercoiled plasmid DNA, which promotes oxidative DNA damage. DNA damage is accompanied by cell death and mutation, as well as associated CVD, neurodegenerative diseases, ageing, and cancer. The NO[•] and O₂^{•-} are the precursors of [•]OH and ONOO⁻, whereas Fe²⁺ and Cu⁺ ions are also known to generate [•]OH by reacting with H₂O₂. Antioxidants impede [•]OH and ONOO⁻-induced DNA damage by scavenging O₂^{•-} and [•]OH.^{27,60-65}

According to Kang *et al.*⁶³ three polyketides, which include cyathuscavins A, B, and C isolated from mycelium culture of *Cyathus stercoreus*, elicited antioxidant activities against DNA damage and thereby protect the supercoiled plasmid DNA from Fe²⁺/H₂O₂-induced breakage. Melatonin, an antioxidant, protected DNA from oxidative damage in a study carried out by Zhao *et al.*⁶⁴ in which each molecule of melatonin trapped approximately two radicals. The inhibitory effect of polyphenols on DNA damage acts by binding to Fe²⁺ ions, as reported by Perron *et al.*⁶⁵ According to Zhao *et al.*⁶⁴ icariin (flavonoid) exhibited antioxidant activity against 2,2'-azobis (2-amidinopropane) dihydrochloride (AAPH)-induced oxidative DNA damage in a dose-dependent manner.

Lipid Peroxidation Inhibition

Lipid peroxidation is the oxidative deterioration of lipids made up of carbon-carbon double bonds present in phospholipids, unsaturated fatty acids, glycolipids, cholesterol esters, and cholesterol. The destructive mechanism of ROS is initiated by electrophilic attack of unsaturated fatty acids that are made up of multiple double bonds and -CH₂- groups (Figure 4).^{7,66} Antioxidants, such as vitamins, flavones, and polyphenols, act directly to quench peroxide radicals to obliterate free radicals' chain reactions. Lipid peroxidation leads to various deleterious health outcomes, such as cancer and atherosclerosis. However, ROS and peroxide radicals are scavenged by antioxidants and thus prevent the presentation of certain pathologic conditions.⁷ Polyphenols from green tea, α-tocopherol, and ascorbic acid acted synergistically to inhibit peroxidation of linoleic acid in SDS micelles.⁶⁷ Likewise, α-tocopherol at μM concentrations inhibited copper-induced peroxidation of liposomal palmitoyllinoleoyl-phosphatidylcholine as reported by Gal *et al.*⁶⁸

MEASUREMENT OF ANTIOXIDANT ACTIVITIES

Antioxidant activities are measured either *in vivo* or *in vitro* using standard experimental methods. Also, standard methods are available for measurement of actions of pro-oxidants. Generally, antioxidant activity is not measured directly but is based on the ability of antioxidants to control the rate of oxidation. There are different established methods that are used for measuring antioxidant capacity.⁶⁹

During measurement of antioxidant activity, it is necessary to take into consideration the source of ROS and target substrate. For instance, a given antioxidant that inhibits oxidative damage to lipids, on the contrary, could promote oxidative damage of another biological molecule.⁷⁰

Most of the methods used in the measurement of antioxidant activities accelerate the oxidation of the substrate in the presence of a promoter under regulated experimental variables such as pH, temperature, concentration of the antioxidant, etc., of the test system. Nevertheless, changes may occur in the oxidation mechanisms when some of the variables are altered. Thus, to obtain an authentic result, it is crucial to estimate the time interval within which the antioxidant activity quantification was carried out.⁶⁹

The analysis of antioxidant capacity is categorised into two main groups with regards to the reaction mechanisms involved in the free radical reduction process. The first and second groups are based on SET and HAT, respectively. The kinetics and secondary reactions that are involved in the methods vary. The SET-based




Figure 1: Action of enzymatic antioxidants on free radicals.³¹

methods detect the ability of a potential antioxidant to transmit chemical species such as metals, carbonyls, and radicals. In addition, the SET-based methods are expressed by a change in colour that occurs because of the reduction of the oxidant by an antioxidant.⁷¹ The HAT-based methods measure the capacity of an antioxidant to deactivate free radicals through a hydrogen atom donation mechanism. The reactions of HAT do not theoretically depend on solvent nature and pH and occur at a very fast rate (less than a few minutes). However, HAT testing is complicated, and significant errors may occur due to the reducing agents present in the samples in conjunction with the antioxidants under study.¹³

The techniques used in the measurement of antioxidant activity must be fast, reproducible, and carried out using small amounts of the compound being analysed and must not be affected by the physical properties of the compound being analyzed. Since the outcomes of *in vitro* assays can be used as a direct indicator of antioxidant activity *in vivo*, a compound that has low antioxidant capacity *in vitro* will also exhibit low antioxidant capability *in vivo*.⁷⁰ Nevertheless, there may be variations in the results obtained depending on the oxidation conditions used and various factors such as temperature, metal catalysts, and the oxygen concentration in the reaction medium that can influence oxidation. The tests employed for the measurement of substrates

or products can also cause variable results depending on their specificity.⁷² The methods used in the measurement of free radical scavenging capacity of antioxidants *in vitro* and *in vivo* are described below.

IN VITRO METHODS

Total Oxyradical Scavenging Capacity Assay

Total oxyradical scavenging capacity assay TOSCA is based on HAT. The TOSCA evaluates antioxidant activity in the gaseous phase and involves the introduction of KMBA to powerful oxidising agents such as ROO[•], •OH, and ONOO⁻ (Figure 5),¹³ which causes the conversion of KMBA to ethylene. The basis of the TOSCA assay is the ability of antioxidants to inhibit the formation of ethylene from KMBA, whereby the antioxidants compete with KMBA for ROS. The capacity of antioxidants is assessed by comparing the extent of ethylene formation with a control reaction using HS-GC. The TOSCA is not suited for a high-performance analysis due to the multiple injections of each sample needed for the estimation of ethylene production.^{13,69} A major limitation of this method is that the concentration of the antioxidants cannot be measured by the percentage inhibition of KMBA oxidation.⁷³




Figure 2: Fe^{2+} binding mechanisms of antioxidants (A) taxifolin, (B) usnic acid, (C) resveratrol and (D) curcumin.⁵¹




Figure 3: The nitration and chlorination of tyrosine.⁷

2,2-Diphenyl-1-picrylhydrazyl (DPPH) Radical Scavenging Assay

The DPPH radical scavenging assay is based on SET, with HAT mechanism as the only marginal reaction pathway.¹³ The DPPH is a stable chromogen radical characterised by a deep purple colour. The basis of the DPPH scavenging assay is the donation of electrons by antioxidants for the neutralisation of DPPH radicals, and the antioxidant activity is measured by the colour change of DPPH at $\lambda_{\max} = 517$ nm.⁷⁴ The DPPH assay is a very simple process that involves only an ultraviolet spectrophotometer or an EPR spectrometer. The EPR spectroscopy directly estimates DPPH radical concentration at sub-micromolecular levels. This assay is majorly dependent on the assumption that the antioxidant electron-donating power (or reducing ability) is directly proportional to the antioxidant activity. Although this method is very simple and does not need special sample treatment, the sensitivity of this assay may be influenced by certain factors such as the type and quantity of the solvent used, the availability as well as the concentration of hydrogen and metal ions, and the viability of DPPH reagent.⁷⁵⁻⁷⁷ The main limitation of the spectrophotometric DPPH assay is the overlapped spectra of compounds that absorb at the same wavelength range as DPPH. A good example is anthocyanins that elicit strong absorption over an equivalent wavelength range as DPPH ($\lambda_{\max} = 500-600$ nm), which interferes with the results and affects their interpretation. To overcome this drawback, the use of ultraviolet spectrophotometry should be substituted with EPR spectroscopy. The EPR spectroscopy is more effective than the classic spectrophotometric method for both highly coloured samples and samples that appear turbid in the chosen solvent.⁷⁸

The DPPH radical possesses an unpaired electron, producing a characteristic and distinguishable EPR spectrum in a magnetic field under microwave spectrum. The peak intensity in the EPR

spectrum is equal to DPPH concentration, and the presence of antioxidants causes its decline with time. The decline in DPPH signal intensity is used as an indicator of antioxidant scavenging capacity against DPPH radical.⁷⁹ Antioxidants scavenging capacity of DPPH radical can also be measured using amperometric detection. The DPPH radical produces a constant electric current at an applied voltage. However, the amperometric signal decreases when DPPH radical concentration drops due to scavenging action by antioxidants. The antioxidants are then quantified using amperometric detection of residual concentration of unreacted DPPH radical.^{80,81} The DPPH radical scavenging assay can also be used in conjunction with other methods for purposes. For instance, DPPH assay can be used in concert with online HPLC to enhance the screening of large quantities of antioxidant samples and thereby decrease the number of antioxidants that will be lost through the purification processes.⁷⁵

Empirically, the methods by Manzocco *et al.*⁸² measure DPPH radical scavenging percentage of sample extract diluted in methanol mixed with DPPH solution. The absorbance is read after 30 min at 517 nm. The percentage inhibition of DPPH radical scavenging in the presence of the sample is calculated (Eqn 14):

$$\% \text{ inhibition of DPPH radical} = \frac{[A_{br} - A_{ar}]}{A_{br}} \times 100 \quad (14)$$

Where A_{br} and A_{ar} are the absorbance before and after the reaction has occurred respectively.

Total Radical-Trapping Antioxidant Parameter

TRAP measures the status of a secondary antioxidant in plasma and functions via HAT. The TRAP is based on the measurement of the rate of uptake of oxygen during a controlled peroxidation reaction following thermal decomposition of ABAP, which is associated with the release of ROO^{\bullet} at a constant rate (Figure

6). This process is carried out by first adding ABAP to human plasma. The parameter being evaluated is the 'delay time' within which oxygen is absorbed in the plasma that is influenced by antioxidants present in the medium. The delay time is estimated through oxygen concentration data in plasma diluted in a buffer solution observed using an electrode. In the absence of ABAP, other free radical initiators, namely, luminol, ABTS, phycoerythrin, and dichlorofluorescein diacetate can be used.⁸³⁻⁸⁵ The results of this method are reported in μmol of ROO^\bullet trapped per litre of plasma.⁶⁹ The major limitation of TRAP is that error may arise during detection of the end point due to instability of the oxygen electrode, which at this point can take up to 2 hr for oxygen to stabilize. However, this problem can be overcome by carrying out electrochemical detection of oxygen using chemiluminescent detection based on the use of luminol and horseradish peroxidase.⁶⁹

The TRAP of diluted samples in phosphate buffer, mixed with bi-distilled water, R-phycoerythrin, and ABAP, is measured according to the methods of Ghiselli *et al.*⁸³ using a luminescence spectrometer. The TRAP is computed using the length of the lag phase due to the sample compared with the standard.

Crocin-Bleaching Assay

The CBA is used to measure the rate at which RO^\bullet generation is inhibited. The reaction mechanism of this assay is based on HAT. The CBA is carried out by evaluating the protective effect elicited by antioxidants on crocin (a carotenoid that gives an intense red colour upon encountering RO^\bullet) (Figure 7). This is made possible using an ultraviolet-visible spectrophotometer for a measure of the reaction kinetics. The absorbance is measured at a wavelength of 440 nm, whereby the relative velocity constants are obtained. The antioxidant capacity of the reference compounds is estimated through the relative velocity constants. The absolute bleaching velocity of crocin is dependent on the kind of radical that attacks the polyene structure of crocin. Crocin shows a high selectivity for RO^\bullet formed from OOH photolysis with RO_2^\bullet formed after the thermolysis of azo initiators.⁸⁶

Oxygen Radical Absorbance Capacity (ORAC) methods

The ORAC methods are centered on the prevention of oxidation initiated by RO_2^\bullet and simultaneously measure the time effect as well as the degree of inhibition. The reaction mechanism of action of this test is based on HAT and utilises the ability of antioxidants to prevent the degradation of fluorescein (fluorescence probe) by promoting the elimination of radicals.⁸⁷ The ORAC starts with the thermal degradation of azo compounds, such as AAPH. This degradation leads to the release of free radicals. The free radicals enhance the degradation of fluorescein. However, the antioxidant

activity promotes the removal of RO_2^\bullet and thereby prevents the degradation of fluorescein. The dilapidation in fluorescence caused by radicals and antioxidants protective effects results in a curve. The area under the fluorescence decrease curve is used to estimate the extent of the antioxidant effect. The standard used in this assay is Trolox. And therefore, the antioxidant activity is measured in terms of Trolox equivalents. The ORAC has been used extensively to measure the antioxidant capacity of vegetables, fruits, supplements, and beverages.^{13,88,89} The ORAC is performed at about pH 7.4, which is adjusted using a phosphate buffer with the antioxidant, AAPH, and fluorescein at 37°C. Fluorescence is observed at 1 min interval for a period of 35 min at an excitation wavelength and emission wavelength of 485 nm and 520 nm, respectively. Furthermore, the modification of this method involves the use of fluorescein as a probe to estimate lipophilic and hydrophilic compounds as well as the total antioxidant capacity of the compound.⁹⁰

According to the method by Prior *et al.*⁹¹ the ORAC assay is performed in polypropylene fluorescence plates. The hydrophilic and lipophilic assays in the presence of AAPH are carried out using Trolox in standard phosphate buffer. The fluorescence is measured at excitation and emission wavelengths of 485 nm and 520 nm, respectively. Wallac Workout 1.5 software is used for the computation of the area under the curve. The results are gotten from the difference of the areas-under-the-decay curves between blank and sample and/or standard (Trolox), expressed as $\mu\text{M TE/g}$ (μM of Trolox equivalents, TE per g dry weight of sample).




Figure 4: Linoleic acid peroxidation mechanism by hydroxyl radical initiation.⁸

IN VIVO METHODS

For all *in vivo* methods, blood or tissues collected from animals such as rats, mice, etc., treated with the samples whose antioxidant efficacy are to be tested, are used for the assay.⁹² The levels of lipid peroxidation, FRAP, GSH, and antioxidant enzymes such as GP_x, GST, SOD, CAT, GGT, and GR are estimated.

Lipid peroxidation

MDA is used as an indicator of lipid peroxidation. MDA is among the end products of lipid peroxidation, and it is generated during oxidative degeneration as a product of free oxygen radicals.⁹² The methods by Ohkawa *et al.*⁹³ for the estimation of lipid peroxidation have been reported elsewhere.^{94,95}

Ferric reducing ability of plasma

Antioxidant efficacy is evaluated through the measurement of the rise in absorbance that occurred by the generation of ferrous ions from FRAP reagent. The absorbance is measured at 593 nm using a spectrophotometer.^{92,96}

Reduced glutathione

GSH is an intracellular reductant that guides the cell from damage by free radicals, peroxides, as well as other toxic substances.⁹⁷ The measurement of GSH in tissue homogenate using Ellman's reagent was described.⁹⁸ The values of the absorbance at $\lambda_{\max}=412$ nm are compared with a standard curve gotten from known GSH.

Glutathione peroxidase activity

Cytosolic GP_x activity is measured by a coupled reaction involving the conversion of NADPH to NADP in the presence of glutathione reductase. The change of absorbance at $\lambda_{\max}=340$ nm of the sample mixture is measured at regular intervals of 1 min for 5 min.⁹⁹ The GP_x activity is expressed as mg of proteins.

Glutathione S-transferase activity

GST activity is measured by the conjugation of CDNB with GSH (Eqn 15) at $\lambda_{\max}=340$ nm at 37°C.^{100,101}



The increase in absorbance is measured for 10 min against a blank solution using a spectrophotometer.




Figure 5: Reaction between KMBA and ROO• to form ethylene.

Superoxide dismutase activity

According to McCord and Fridovich,¹⁰² SOD activity is measured in erythrocyte lysate prepared from the 5% erythrocyte suspension. The sample mixture consists of Tris-HCl buffer (pH 8.2), EDTA, and pyrogallol. The increase in absorbance is measured at $\lambda_{\max}=420$ nm for 3 min against a blank solution using a spectrophotometer. One unit of SOD activity is equivalent to the auto-oxidation rate of pyrogallol with regards to the change in absorbance/min. SOD enzyme activity is expressed as units/mg protein.

Catalase activity

CAT activity of erythrocyte lysate is measured in phosphate buffer (pH 7.0). The enzyme activity is evaluated using the molar extinction coefficient of H₂O₂ ($\Sigma=43.6$ M cm⁻¹). One unit CAT activity is equivalent to 1 mmol of H₂O₂ consumed per min. Catalase activity is expressed as units per milligram of protein.¹⁰³

γ -Glutamyl transpeptidase activity

The γ -GGT activity is measured according to the methods previously described.¹⁰⁴ The serum sample is pre-mixed with MgCl₂, glycylglycine and γ -Glutamyl-p-nitroanilide in tris buffer (free base) (pH 8.2) incubated for 1 min at 37°C. The absorbance is measured at regular interval of 1 min for 5 min at $\lambda_{\max}=405$ nm.

Glutathione reductase activity

The oxidation rate of NADPH by GSSG at 30°C is used as a standard method for the measurement of GR activity. The reaction system is made up of NADPH, GSSG, EDTA and sodium phosphate buffer (pH 7.6). The Unit of GR activity is derived from the oxidation of NADPH/min. The specific activity of GR is stated as Unit per mg of protein.¹⁰⁵

OXIDATIVE STRESS AND PATHOLOGY

Pathological conditions such as CVD, DM, tissue inflammation, sickle cell anaemia, cancer, cataracts, AD, autism, and ageing are provoked by increased free radical generation in conjunction with alteration in system antioxidant defence capabilities. Thus, indicating the contributive role of ROS in pathological on set, escalation, and complications. Increasing evidence in both experimental and clinical studies suggests that oxidative stress contributes largely to the pathogenesis of various disease conditions.¹⁰⁶⁻¹⁰⁸

It's conventional to integrate antioxidant formulations into therapeutic interventions, whereby natural antioxidants administered through dietary supplementation are recommended for a holistic approach to the management and prevention of oxidative stress-induced pathological conditions and complications. An earlier proposed unique therapeutic strategy, whereby overexpression of enzymatic antioxidants was provoked in a mutant mice model in a tissue-specific manner, could serve as a novel alleviative measure against the development of pathological conditions associated with oxidative stress.^{109,110} Similar novel approaches have been adduced against human inflammatory pathways associated with diabetic complications.¹¹¹ Specifically, one of such approaches is the activation of transcription nuclear factor, Nrf2, which in turn induces several enzymatic antioxidants and detoxification genes against oxidative stress-induced pathology such as lung cancer.^{110,112}

Cardiovascular Disease

The ROS plays a dual role in cardiovascular physiopathology. Low levels of ROS provide significant benefits to the cardiovascular system, whereby they exhibit pro-angiogenesis, anti-atherosclerotic, and endogenous cardio-protective effects.¹¹³ On the contrary, high levels of ROS engender deleterious actions, through oxidative stress, in the cardiovascular system by provoking CVD, such as atherosclerosis, heart failure, endothelial dysfunction, arrhythmias, and myocardial ischaemia/reperfusion.^{113,114} In addition, oxidative stress causes alteration in gene expression in CVD.¹¹⁵ At high levels of ROS, one of the foremost occurrences is oxidative modification of LDL in atherogenesis and other CVD associated with endothelial dysfunction.^{116,117} Furthermore, LDL and cell membranes are composed of large amounts of phospholipids, which are sensitive to oxidative modification. Through receptor-mediated or receptor-independent pathways, oxidised phospholipids stimulate endothelial cells, enhance the expression of endothelium adhesion molecules, attract monocytes, elicit endothelium cytotoxic effects, and promote the activities of pro-inflammatory genes and cellular growth factors. All these events elicit aggregation of platelets, expression of metalloproteinase, and promote thrombogenesis.¹¹⁸

Generally, endothelial dysfunction initiated by excessive ROS generation overwhelms antioxidant defence systems, causing damage to endothelium-mediated vasodilation, impaired vascular reactivity, and vasospasm, as well as increased expression

of chemotactic and adhesive molecules. And thereby, enhances activation of platelet and thrombus generation, increased permeability of the endothelium, leucocyte adhesion, as well as monocyte movement into the vascular wall. The regeneration of endothelial cells, accompanied by the proliferation and migration of smooth muscle cells, is impaired, and in the process, engenders vascular damage (Figure 8).¹¹⁹⁻¹²¹

The major risk factors of CVD, namely, hypercholesterolaemia and hypertension, promote ROS generation, which eventually leads to oxidative stress.¹²² Higashi *et al.*¹²³ investigated the level of 8-OHdG in urine as well as serum MDA-modified LDL as markers of oxidative stress in patients suffering from reno-vascular hypertension. Higher levels of these oxidative stress indicators were observed in hypertensive patients in comparison with the non-hypertensive control group. Another study by Lip *et al.*¹²⁴ also reported elevation of biomarkers of oxidative stress in hypertensive individuals.

Oxidative stress induces the rupture of atherosclerotic plaque and eventually causes thrombosis by stimulating the expression of matrix metalloproteinase.¹²⁵ The action of NF- κ B is upregulated by oxidised LDL in atherosclerosis.¹¹⁶ The study carried out by Guzik *et al.*¹²⁶ showed a correlation between endothelial dysfunction and elevated vascular $O_2^{\cdot-}$ generation in human atherosclerosis. Azarsiz *et al.*¹²⁷ reported the relationship between oxidative stress and atherosclerosis, in which they noted increased erythrocyte levels of TBARS with corresponding severity of blockage of the arteries.

Because cardiac myocytes possess higher amounts of mitochondria in comparison with other cells, they consume relatively higher levels of oxygen to generate energy in the form of ATP. The relatively high level of non-neutralised ROS generated as a byproduct of tissue metabolism causes cardiac injury in the myocytes, with concomitant oxidation of essential proteins required for the excitation-contraction process and mitigation of the bioactivity of NO.¹²⁸

In myocardial ischaemia, there is an imbalance in mitochondrial electron transport with concomitant ATP level reduction, acidosis, and depolarisation of the mitochondria, intracellular Ca^{2+} overload, and apoptosis.¹²⁹ Oxidative stress generated in the mitochondria causes impaired mitochondrial DNA and presentation of CVD. Furthermore, in myocardial ischaemia, free radical generation is enhanced in the cardiac tissue by hypoxia




Figure 6: Peroxyl radical formation from 2,2'-azobis(2-amidopropane).




Figure 7: Molecular structure of crocin.

and re-oxygenation.¹¹⁵ The ROS generated via re-oxygenation initiates direct oxidative damage to cellular components and indirect impairment by provoking localised inflammation.¹³⁰ The increased ROS generation in heart failure is dependent on the elevated activity of xanthine oxidoreductase and NADPH oxidase. High levels of ROS are accompanied by long-term endoplasmic reticulum stress as well as mitochondrial-derived oxidative stress in cardiometabolic disorders.¹¹⁴

Diabetes Mellitus

Experimental and clinical studies suggest that oxidative stress contributes significantly to the pathogenesis and development of DM complications. Nevertheless, the mechanism through which oxidative stress contributes to and enhances DM complications is not fully understood and needs to be fully clarified.

Hyperglycemia activates free radical generation and alters the endogenous antioxidant defence system in diabetic individuals.¹³¹ There are eight main mechanisms that lead to free radical generation through uncontrolled hyperglycemia, and they include an increase in the flux of the polyol pathway, glucose autoxidation, an increase in the rate of glycolysis, increased AGEs generation, increased expression of the receptor for AGEs as well as its activating ligands, non-enzymatic glycation of proteins, NAD(P)H oxidase activation through the protein kinase C-dependent pathway, and increased activity of the hexosamine pathway.^{132-136,110} The formation of oxidants in DM and the concomitant oxidative stress leading to oxidative cellular damage is described in Figure 9.

RAGE: Receptor for AGEs

Furthermore, the eight major pathways/mechanisms contributing to the pathogenesis of diabetic complications are induced by excessive $O_2^{\cdot-}$ generation via hyperglycaemic intracellular setting (Figure 10).^{135,138-141} Specifically, there is an increase in the level of pyruvate produced from glucose catabolism with high intracellular glucose concentration, thereby making more pyruvate available for oxidation via the TCA cycle, which increases the flow of NADH and $FADH_2$ (electron donors) into the electron transport chain. This process causes a continuous increase in voltage gradient across the mitochondrial membrane until a critical threshold is reached. At the threshold point, the

transfer of electrons in the mitochondrial electron transport chain causes $O_2^{\cdot-}$ generation through the leakage and transfer of one electron to molecular oxygen (Figure 10).¹⁴¹

In addition, hyperglycemia also activates GAPDH, which eventually leads to a high accumulation of glycolytic intermediates that are upstream of this enzyme.¹⁴² The highly accumulated glycolytic intermediate, glyceraldehyde-3-phosphate, activates the PKC pathway (because diacylglycerol, which is the activator of PKC, is formed from glyceraldehyde-3-phosphate) as well as the pathway that leads to AGE generation (the chief intracellular precursor of AGE, methylglyoxal, is produced from glyceraldehyde-3-phosphate). The flux of the hexosamine pathway is also increased through the accumulation of another glycolytic intermediate, fructose-6-phosphate, which is upstream of GAPDH. Also, an increase in the level of glucose through the inhibition of GAPDH by hyperglycemia increases the polyol pathway flux (Figure 10).^{141,143-146}

A study carried out by Davison *et al.*¹⁴⁷ suggested that a high level of oxidants in type I DM male individuals was because of the autoxidation of glucose and/or an increase in the rate of oxidation of the major lipid-soluble antioxidant, α -tocopherol, due to low exercise. Hyperglycemia was observed to induce overproduction of oxygen free radicals in children with chronic insulin-dependent DM in a study carried out by Ramakrishna and Jaiikhani.¹⁴⁸ According to the study carried out by Godin *et al.*¹⁴⁹ using streptozotocin-induced diabetic rats, it was reported that elevated and uncontrolled oxidative activities promoted the pathogenesis and complications that are associated with chronic diabetic conditions. The increase in susceptibility of erythrocyte lipids to oxidation was directly proportional to the severity of diabetic complications. Additionally, certain changes, such as elevation in GR activity, reduced vulnerability to decreased oxidative glutathione (which is associated with hyperglycemia), and elevation in malondialdehyde generation (which is an indirect measure of lipid peroxidation), were observed in the diabetic state. Yakubu *et al.*¹⁰⁷ were of the view that the pathogenesis of glucose-induced vasculopathy correlated with PKC-coupled oxygen free radical generation, which suppressed NO^{\cdot} generation and NO^{\cdot} -dependent vascular relaxation. The previous report by Ohkuwa *et al.*¹⁵⁰ suggested that $\cdot OH$ generation was dependent on the amount of streptozotocin injected into experimental rats.

And therefore, they concluded that $\cdot\text{OH}$ generation is involved in the pathogenesis of DM.

The production of $\text{O}_2^{\cdot-}$ was reported by Ceolotto *et al.*¹⁵¹ to be induced by insulin through an NAD(P)H-dependent mechanism, which involved the activation of PI-3'-kinase and activation of proliferative ERK-1 and -2 dependent pathways. An elevation in oxidative stress was observed to be accompanied with a marked reduction in the level of GSH as well as alterations in enzymatic antioxidant activities such as catalase, GPx, and Cu-Zn-superoxide dismutase in the heart, liver, and kidneys of streptozotocin-induced diabetic rats.¹⁰⁶ Similar changes in the enzymatic antioxidants in rat brain were also reported.¹⁰⁸

Cancer

Oxidative stress in cancerous cells arises when ROS generation overwhelms the antioxidant defence system, leading to DNA, protein, and/or lipid damage; chromosomal instability; genetic mutation; as well as cell growth modulation, which eventually results in cancer (Figure 11).¹⁵²

Free radicals cause chromosomal defects as well as the activation of oncogenes, which eventually leads to the emergence and progression of cancer.¹⁵³ High levels of free radicals have been reported to cause cell damage and apoptosis. Free radicals and DNA reactions have been linked to being associated with various forms of cancer, leading to mutations that have a deleterious effect on the cell cycle and eventually result in neoplasia.^{154,155}

Excessive ROS generation promotes the proliferation of cancer cells, metastatic potential, as well as invasiveness and poor prognosis.¹⁵⁶ ROS promotes the migration of cancer cells by supporting matrix degradation, which facilitates contact among cells, cytoskeleton remodelling, as well as regulates gene expression and invadopodia formation.¹⁵⁴ For example, mitochondria-derived ROS influences initial extracellular matrix contact; NOX-derived ROS are associated with invadopodia formation, and an increase in the level of ROS in the cytosol contributes to cytoskeleton remodelling.¹⁵⁷

The ultraviolet component of sunlight, within wavelengths of 320-400 nm, has the potency to initiate oxidative stress in tissues. And continuous exposure of human skin to ultraviolet radiation is accompanied by skin carcinogenesis as well as photoaging through DNA damage. The physiological concentration of ultraviolet components of sunlight affects the expression of certain genes, such as nuclear oncogenes, collagenase, and heme oxygenase 1, whose effects can be significantly increased through the removal of intracellular GSH or by prolonging the half-life of molecular oxygen.^{158,159} The study by Sander *et al.*¹⁶⁰ reported that oxidative stress contributed to the pathogenesis of human skin cancers and suggested that in non-melanoma skin cancer, the reduction in antioxidant defence system induced by exposure to chronic ultraviolet radiation may promote multistep

carcinogenesis, while melanoma cells exhibit increased oxidative stress that can initiate tissue damage and promote metastasis.

The $\cdot\text{OH}$ -attach to DNA reacts with deoxyGTP in the nucleotide pool to generate 8-OHdG, which causes mutation in mammalian and bacterial cells (Figure 12).¹⁵²

The 8-OHdG also induces the transformation of G-C pairs to T-A pairs during the process of DNA replication, which eventually leads to the emergence of cancer-inducing mutagenesis.¹⁶¹ Thus, 8-OHdG can serve as a biomarker for the detection of the presence of free radicals during DNA mutagenesis.¹⁶² Many studies have reported high levels of 8-OHdG in various human cancers¹⁶³⁻¹⁶⁶ as well as in animal models of tumours.^{167,168} Furthermore, ROS accumulation is associated with an increased proliferation of cancer cells.¹¹⁴

There is continuous oxidative stress in cancer cells, which is enhanced by mitochondrial dysfunction as well as metabolic changes. Under normal conditions, high ROS levels induce cell death, but this does not occur in cancer cells due to their ability to activate numerous oncogenes, which in turn activates the expression of NRF2. The NRF2 is the primary regulator of cell survival that increases cancer progression by defending cancer cells from the damages caused by ROS on DNA.^{114,169} The ROS are implicated in the progression of cancer through the promotion of cyclin D1 expression, ERK, and the phosphorylation of JNK.¹⁵⁹ The investigation carried out by Aikawa *et al.*¹⁷⁰ reported the role of ROS in the activation of ERK in cultured cardiac myocytes of neonatal rats.

Alzheimer's Disease

AD is one of the most encountered neurodegenerative diseases that lead to dementia among the elderly. The AD is characterised by gradual loss of memory as well as other cognitive functions, which finally leads to the complete incapacity and death of the affected person within a period of 3 to 9 years following diagnosis.¹⁷¹ The presence of senile plaques, NFTs, and neuronal cell death are the main pathological characteristics of AD brains.¹⁷²

Cytochrome c oxidase deficiency is the major and common impairment in the mitochondrial ETC in AD, which results in increased ROS generation.¹⁷³ The ROS triggers the accumulation of A β protein, which leads to lysosome membrane degradation as well as neuronal death.¹⁷⁴ Furthermore, ROS suppresses PP2A, leading to the activation of GSK 3 β and probably the hyperphosphorylation of tau and neurofibrillary lesions.^{175,176} Elevated ROS initiates low mitochondrial membrane potential ($\Delta\Psi_m$) and ATP generation by unfavourable mitochondrial energy stores, disrupting energy metabolism, as well as compromised dynamics and mitophagy. These events further trigger caspase activity and apoptosis (Figure 13).¹⁷⁷ Studies have shown that mitochondrial dysfunction is a major factor that is associated with the pathogenesis of AD.^{178,179} Many mitochondrial and

metabolic abnormalities occurred in the hippocampal neurones in AD compared with age-matched controls.^{178,179} Morphometric analysis of biopsies from AD brains reported a substantial reduction in mitochondria, whereas mitochondrial DNA and protein located in the cytoplasm and vacuoles associated with lipofuscin showed significant increase.¹⁷⁸⁻¹⁸⁰ These mitochondrial abnormalities were followed by oxidative damage marked by raised levels of 8-OHdG and nitrotyrosine, which showed that mitochondrial impairment occurred during AD progression.¹⁷⁸

Many hypotheses are in support of oxidative stress as one of the factors involved in the pathogenesis of AD. However, the mechanism through which the redox balance is altered, and the sources of free radicals have not been established.¹⁸¹

Studies have also reported possible correlation between oxidative stress and A β protein-induced toxicity. Increased H₂O₂ and NO \cdot generation coupled with enhanced oxidative modifications of proteins and lipids were directly proportional to the age-associated A β protein accumulation in various AD transgenic mouse models carrying mutants of APP and PS-1, which suggested that A β protein promoted oxidative stress.¹⁸²⁻¹⁸⁵ In the study carried out by DeFelice *et al.*¹⁸⁶ using hippocampal neuronal cell cultures, ROS activated by soluble A β protein oligomers triggered NMDA receptor, which was accompanied by a sudden rise in the levels of neuronal calcium. The report therefore suggested a possible role of soluble A β protein oligomers as proximal neurotoxins and the

active role of oxidative stress in synaptic damage and neuronal loss initiated by soluble A β protein oligomers.

Alterations in metal homeostasis have been shown to relate to AD. Abnormal levels of Cu, Zn, and Fe were reported to occur in the hippocampus and amygdala regions with severe histopathologic alterations in AD.¹⁸⁷ Transition metals such as Cu, Zn, and Fe are known to play important roles in neural functions. However, metal levels and transportation are stringently regulated because abnormal metal homeostasis can lead to neurotoxic free radical generation. For instance, Fe or Cu at very high levels can directly interact with oxygen, generating O₂ \cdot^- , H₂O₂, and \cdot OH and possibly resulting in oxidative stress as well as a cascade of biochemical alterations that eventually leads to neuronal cell death.¹⁸⁸ Furthermore, studies have also reported the presence of these transition metals within the amyloid deposits of patients suffering from AD and transgenic mouse models.^{189,190}

Parkinson's Disease

PD is one of the major neurological disorders that are very prevalent among individuals above 65 years of age.¹⁹¹ The PD is a progressive neurodegenerative disorder that affects, primarily, dopamine neurones that originate from the midbrain and extend to the putamen and caudate regions (the striatum), which are regions that are responsible for the control of motor movements.¹⁸¹ The PD is characterised by loss of dopaminergic neurones in the SNpc. Oxidative stress promotes dopaminergic neurotoxicity. Complex I deficiencies of the respiratory chain contribute




Figure 8: Endothelial dysfunction leading to vascular damage initiated by oxidative stress.¹²¹




Figure 9: Oxidant formation in diabetes mellitus and the concomitant oxidative stress leading to oxidative cellular damage.^{110,137}

significantly to most of the adverse neuronal degeneration in PD. Dopamine, neurotoxins, and mutations of genes in PD-associated proteins play a major role in mitochondrial dysfunction, which leads to ROS generation (Figure 14).¹⁹²

The metabolism of dopamine, which is catalysed by monoamine oxidase, is enhanced in PD, and excess H₂O₂ is generated in the process (Figure 14). The polymerisation of the auto-oxidative products of dopamine can possibly initiate the production of the characteristic pigmentation of the substantia nigra. These free radicals generated might be responsible for the loss of dopaminergic neurons.¹⁸¹

Mitochondrial dysfunction enhances the formation of ROS in PD.¹⁹³ Oxidative phosphorylation supplies most of the energy required to trigger neural activity in which the mitochondria use their structure, enzymes, and energy generated through the oxidation of nutrients to produce ATP.¹⁹⁴ However, this metabolic pathway that occurs in the mitochondria is the major source of O₂^{•-} and H₂O₂.¹⁹²

Complex I deficiencies of the respiratory chain contribute significantly to unfavourable neural apoptosis and are seen as one of the major sources of ROS in PD. The inhibition of complex I results in increased ROS generation. Reports showed a decreased activity of complex I in the SNpc of sporadic PD patients.^{107,195,196} Furthermore, studies have reported mitochondrial complex I

deficiency in various brain regions, skeletal muscle, fibroblasts, lymphocytes, and blood platelets of PD patients.¹⁹⁷⁻²⁰²

Other evidence that supports the relationship between mitochondrial dysfunction and oxidative stress as well as dopamine cell damage arises from discoveries that mutations in genes of proteins such as parkin, PINK, α -syn, or DJ-1 are associated with familial forms of PD.^{203,204} Generally, these findings suggested that mutations in these genes altered the function and integrity of the mitochondria and triggered oxidative stress (Figure 14).²⁰⁵

According to Wang *et al.*²⁰⁶ PINK-1 plays a major role in the maintenance of normal morphology of mitochondrial SNpc dopamine neurones and eliciting a neuroprotective effect by suppressing ROS generation. Studies carried out using animal models showed that deficiency of PINK-1 resulted in impaired mitochondrial morphology, decreased activity of complex I, loss of SNpc dopamine neurones, and increased oxidative stress.²⁰⁷⁻²⁰⁹ The accumulation of α -syn as intracellular filamentous aggregates serves as a pathological indicator of sporadic and familial PD.²¹⁰ Also, accumulation of α -syn wild-type in dopamine neurones decreased mitochondrial complex I activity with increased ROS generation, and cell death ensues.²¹¹ Dendritic mitochondrial oxidative stress was increased by α -syn inclusions in dopamine neurones in the study carried out by Dryanovski *et al.*²¹² Oxidative stress has been reported to promote influx, assemblage, and oligomerisation of extracellular α -syn in oligodendrocytes and

promote posttranslational modifications of α -syn, which may lead to increased dopamine toxicity.^{213,214}

Aging

Ageing is a gradual and complex event that is associated with progressive and permanent deterioration of the cells, tissues, organs, and the whole organism itself and can be referred to as pathological conditions that have effects on the individual's quality of life.²¹⁵

According to the free radical theory of ageing, which was postulated about 60 years ago, ROS is recognised as the main cause of impairments of the cells and responsible for ageing and ageing-associated degenerative diseases.²¹⁶ Although there are many theories of ageing today, oxidative stress remains a causative factor.²¹⁷ In the study by Kumsta *et al.*²¹⁸ *Caenorhabditis elegans* was subjected to oxidative stress via sub-lethal short treatment of H₂O₂ stress. The report showed that most of the worms exhibited symptoms of severe but reversible behavioural changes that are evidence of common age-related changes, such as reduction in body movement, decreased pharyngeal pumping, low reproduction rate, morphological changes, and decline in metabolic activity. These findings were in accordance with the Harman free radical theory of ageing.²¹⁸

The mechanism through which oxidative stress elicits cell aging process is as follows: (1) the metabolism of the cell causes the formation of ROS, which is followed by oxidative damage; (2) the macromolecule mostly affected by oxidative stress are proteins as they undergo various modifications in order to prevent being

properly degraded and recycled by the proteasome and protein function is altered in the process; (3) cytoskeletal proteins are also affected by oxidative stress, leading to structural impairment and signaling alterations; (4) oxidative stress further disrupts energy generation in the mitochondria; (5) the peroxisomes are affected by oxidative stress, which is accompanied by alteration of normal metabolic functioning; (6) the cellular membrane is also affected. (7) Finally, all these processes disrupt the cell's transcriptional activity and cause altered gene expression, which leads to the cell ageing process with concomitant degenerative diseases (Figure 15).²¹⁵

Sickle Cell Disease

Oxidative stress contributes to multiple pathophysiologic mechanisms, such as endothelial damage,^{219,220} hypercoagulability,²²¹ reduced NO• bioavailability,²²² and increased haemolysis²²³ in SCD, which eventually trigger SCD-related vaso-occlusion and chronic organ damage (Figure 16).²²⁴⁻²²⁶ A cross-sectional study carried out by Atiku *et al.*²²⁷ to compare the levels of oxidative stress biomarkers in plasma samples from SCD and healthy control participants showed elevated levels of the ROS biomarker malondialdehyde in SCD individuals in comparison with the healthy control subjects. Higher levels of •OH and NO• were also identified in the plasma of SCD children than those of control groups by Biswal *et al.*²²⁸

In SCD, the endothelium is the main target of oxidative stress due to its nearness to cell-free haemoglobin and regions of ischaemia-reperfusion injury. Many oxidative stress-related mechanisms are involved in endothelial damage, which contributes significantly to SCD-related chronic organ complications such as pulmonary hypertension, cerebral vasculopathy, and retinopathy (Figure 16).^{220,225,226} The ROS incur direct damage on endothelial cells through oxidative reactions with (membrane) lipids, peptides, and nucleic acids.^{226,229} The oxidative events lead to abnormal enlargement of the endothelial cells, resulting in their separation from the underlying basement membrane and exposure of sub-endothelial structures as well as proteins and TF, which supports hyper-coagulation associated with SCD.²³⁰ Furthermore, by inducing NF- κ B, ROS triggers gene expression and pro-inflammatory cytokine generation, such as IL-1, IL-6, and TNF- α , as well as adhesion molecules, namely, VCAM-1 and ICAM-1,²³¹ which eventually enhances a pro-inflammatory state. The enhanced expression of endothelial adhesion molecules might play a role in vaso-occlusive events in SCD through the attachment of sickle erythrocytes, leukocytes, as well as platelets to the endothelium.²³²

Decreased bioavailability of NO• is a major feature of SCD, and this component is essential for vaso-dilatative, anti-inflammatory, and antithrombotic functions.²³³⁻²³⁶ Oxidative stress plays a major role in the reduced bioavailability of NO• in SCD, which takes place as follows: the O₂^{•-} binds to NO•, leading to the formation




Figure 10: High levels of superoxide generation in the mitochondria under hyperglycemic conditions leading to the activation of four major pathways associated with hyperglycemic damage through the suppression of GAPDH.¹⁴¹

NADH: Nicotinamide adenine dinucleotide; NADPH: Nicotinamide adenine dinucleotide phosphate; AGE: Advanced glycation end-product; GFAT: Glutamine: fructose-6-phosphate amidotransferase; GAPDH: Glyceraldehyde-3 phosphate dehydrogenase; PARP: Poly-ADP-ribose polymerase; DAG: Diacylglycerol; UDP-GlcNAc: UDP-N-acetylglucosamine




Figure 11: The role of reactive oxygen species in the development of human cancer.¹⁵²




Figure 12: 8-OHdG formation by hydroxyl radicals.

of the potent oxidant ONOO^- and its conjugate, ONOOH , and thereby further reduces the bioavailability of NO^* and aggravates oxidative stress.²³⁷ Oxidative stress also reduces NO^* bioavailability by contributing to the formation of the eNOS inhibitor ADMA, which is involved in SCD-related pulmonary hypertension and early death.²²²

Increased intra- and extra-erythrocytic oxidative stress enhances intravascular haemolysis by triggering lipid peroxidation as well as membrane instability.²³⁸ High OOH levels cause the degradation and deformity of the erythrocyte membrane.²³⁹ Oxidative stress also plays a role in Ca^{2+} -dependent scramblase-mediated disruption of normal asymmetrical distribution of membrane phospholipids, which causes phosphatidylserine to be exposed at the outer surface of sickle erythrocytes.²⁴⁰ Sickle erythrocytes whose phosphatidylserines are exposed are hydrolysed by secretory phospholipase A2, leading to increased intravascular haemolysis.²⁴¹

Chronic Respiratory Diseases

CRDs refer to a group of chronic diseases that affect the airways and connecting regions of the lungs. Some of the most common

CRDs that are associated with oxidative stress include bronchial asthma, COPD, pulmonary fibrosis, and lung cancer.²⁴² The sources and pathophysiological roles of ROS in CRDs are shown in Figure 17.

Bronchial asthma is a chronic airway disease that is associated with airway eosinophilia, goblet cell hyperplasia accompanied by mucus hypersecretion, and hyper-responsiveness to inhaled allergens and non-specific stimuli, which promote increased vascular permeability leading to plasma exudation.²⁴⁴⁻²⁴⁶ Raised levels of ROS play a significant role in the contraction of airway smooth muscle, alteration in normal functioning of β -adrenergic receptor, reduction in the amount of the epithelial cilia and its function, increased generation of mucus, impairment in the release of inflammatory mediators, inflammatory cells influx, as well as an increase in vascular permeability.²⁴⁷ Additionally, raised levels of ROS initiate bronchial hyperreactivity associated with asthma. Empirical studies using animal models reported that ROS play a significant role in airway hyper-responsiveness by inducing an increase in vagal tone due to the impairment of oxidant-sensitive β -adrenergic receptors and reduction of mucociliary clearance.^{248,249} The study by Henricks and Nijkamp,²⁴⁷




Figure 13: Impairments induced by reactive oxygen species in Alzheimer's disease.¹⁷⁷




Figure 14: Different pathways and their dysfunctions resulting from dopamine metabolism and genetic modifications in Parkinson's disease-related genes and their relationship with oxidative stress.¹⁹²

reported that ROS are involved in endothelial barrier dysfunction and increased permeability to fluid, macromolecules, and inflammatory cells. Previous studies on alveolar macrophages of asthmatic subjects reported higher generation of $O_2^{\cdot-}$ as well as other ROS when compared with those of the healthy controls.^{250,251} More ROS was produced from the eosinophils, neutrophils, and alveolar macrophages of asthmatic patients than those of the normal control subjects.^{252,253}

A COPD is typically associated with progressive, irreversible limitation in the expiratory airflow as well as abnormal inflammation of the lung, which is characterised by chronic inflammation of the airway's parenchyma and pulmonary vasculature, in conjunction with elevated levels of neutrophils,

macrophages, and T lymphocytes (especially CD8+).²⁵⁴⁻²⁵⁶ The pathophysiology of oxidative stress in COPD is complex.




Figure 15: Impairments triggered by oxidative stress that lead to cell aging and age-related diseases.²¹⁵

Persistent oxidative stress was observed in the lungs of COPD




Figure 16: Pathophysiological effect of oxidative stress on sickle cell disease complications.²²⁶




Figure 17: Sources and overall pathophysiological role of reactive oxygen species in chronic respiratory diseases.²⁴³

patients with higher levels of pro-inflammatory cytokine levels, CD4 and CD8 cells, and proteases, as well as increased apoptosis and senescence.²⁵⁷⁻²⁶³ Oxidative stress-activating mechanisms are involved in COPD initiation and progression.²⁶⁴ Oxidative stress biomarkers such as H₂O₂ and NO were seen to be present in the epithelial lining fluid, urine, and breath of the smokers of cigarettes and patients suffering from COPD.^{254,165-167} An increase in the generation of O₂^{•-} and upregulation of adhesion molecules were seen in the circulating neutrophils of patients suffering from COPD.^{268,269}

Pulmonary fibrosis is the outcome of a diverse group of lung impairments. There are suggestions that fibrotic stimuli of unknown origin initiate a state of disequilibrium between the generation of oxidants and antioxidant protection, leading to ROS accumulation.²⁷⁰ Oxidants may contribute to the progress of pulmonary fibrosis through the generation of cytokines as well as growth factors such as TGF-β, which is a major regulator of aberrant repair mechanisms that are characteristic of various fibrotic diseases such as IPF. TGF-β reduces the rate of generation of the natural cellular antioxidant by ameliorating the expression of both CAT and mitochondrial SOD, which further encourages the production of ROS through the activation of NADPH oxidases and/or mitochondrial dysfunction.^{157,271-274} Studies on human subjects suffering from IPF showed increased levels of oxidised proteins.²⁷⁵⁻²⁷⁷ The protective efficacy of antioxidant enzyme systems against lung fibrosis has been reported.^{278,279} The study by Rahman *et al.*²⁸⁰ reported lower antioxidant efficacy in IPF subjects in comparison to healthy subjects.

The ROS have been reported to induce oncogenes such as Jun and Fos. Overexpression of Jun is significantly associated with the occurrence of lung cancer.^{281,283} For instance, in lung cancers, p53, which is involved in the generation of ROS, undergoes mutation, and becomes defective in the activation of apoptosis. The p53 accumulates in the cytoplasm and functions as an oncogene after the mutation.²⁸³

CONCLUSION

The roles of free radicals in pathogenesis are irrefutable and well established. And therefore, the neutralisation and/or scavenging of intracellular free radicals are sustainable therapeutic strategies for the prevention and amelioration of oxidative stress and associated pathological conditions and their complications. Antioxidants prevent cellular damage caused by free radicals by inhibiting the generation and scavenging of radicals or promoting their degradation. Some actions of free radicals impair cellular activities that elicit chronic diseases such as CVD, DM, cancer, AD, PD, and aging. The proposed therapeutic strategy, whereby overexpression of enzymatic antioxidants was provoked in a mutant mice model in a tissue-specific manner, could serve as a novel alleviative measure against the development of pathological conditions associated with oxidative stress.

ACKNOWLEDGEMENT

The authors express their gratitude for the administrative support received from the Secretary and other staff members of the Department of Biochemistry at Imo State University, Owerri, who were immensely helpful during the data collection process.

CONFLICT OF INTEREST

The authors declare that there is no conflict of interest.

ABBREVIATIONS

AAPH: 2,2'-azobis (2-amidinopropane) dihydrochloride; **ABAP:** 2,2'-azobis-(2-amidopropane); **AD:** Alzheimer's disease; **ADMA:** Asymmetric dimethylarginine; **AGEs:** Advanced glycation end products; **AscH:** Ascorbic acid; **BHA:** Butylated hydroxyl anisole; **BHT:** Butylated hydroxyl toluene; **CAT:** Catalase; **CBA:** Crocin-bleaching assay; **-CH₂-:** Methylene; **Cl⁻:** Chloride ion; **CO₃^{•-}:** Carbonate radical; **COPD:** Chronic obstructive pulmonary disease; **CRDs:** Chronic respiratory diseases; **Cu:** Copper; **CVD:** Cardiovascular diseases; **DAG:** Diacylglycerol; **DPPH:** 2,2-Diphenyl-1-picrylhydrazyl; **eNOS:** NO[•] synthase; **ERK:** Extracellular signal-regulated kinase; **ETC:** Electron transport chain; **Fe:** Iron; **GAPDH:** Glyceraldehyde-3-phosphate dehydrogenase; **GFAT:** Glutamine: fructose-6-phosphate amidotransferase; **GPx:** Glutathione peroxidase; **GR:** Glutathione reductase; **GSH:** Glutathione; **GSK:** Glycogen synthase kinase; **GSSG:** Glutathione disulphide; **GST:** Glutathione S-transferase; **H₂O₂:** Hydrogen peroxide; **HAT:** Hydrogen atom transfer; **HMWA:** High molecular weight antioxidants; **HOCl:** Hypochlorous; **HS-GC:** Headspace gas chromatography; **ICAM-1:** Intercellular adhesion molecule-1; **IPF:** Idiopathic pulmonary fibrosis; **JNK:** JUN N-terminal kinase; **KMBA:** α-keto-γ-methylthiobutyric acid; **LDL:** Low-density lipoprotein; **LMWA:** Molecular weight antioxidants; **LOOH:** Lipid hydroperoxide; **MDA:** Malondialdehyde; **Mn:** Manganese; **Mn-SOD:** Manganese-superoxide dismutase; **MPO:** Myeloperoxidase; **NADH:** Nicotinamide adenine dinucleotide; **NADPH:** Nicotinamide adenine dinucleotide phosphate; **NDGA:** Nor-dihydroguaiaretic acid; **NFTs:** Neurofibrillary tangles; **NF-κB:** Nuclear factor-κB; **NMDA:** N-methyl-D-aspartate; **•NO₂:** Nitrogen dioxide radical; **NO[•]:** Nitric oxide; **NO₂Cl:** Nitryl chloride; **Nrf2:** Nuclear factor-erythroid 2-related factor 2; **•OH:** Hydroxyl radical; **•OOH:** Hydroperoxide; **¹O₂:** Singlet oxygen; **O₂^{•-}:** Superoxide anion radical; **ONOO[•]:** Peroxynitrite; **ONOOCO₂^{•-}:** Nitrosoperoxycarbonate; **ONOOH:** Peroxynitrous acid; **ORAC:** Oxygen radical absorbance capacity; **PARP:** Poly-ADP-ribose polymerase; **PD:** Parkinson's disease; **PG:** Propyl gallate; **PKC:** Protein kinase C; **•PP2A:** Phosphatase 2A; **RAGE:** Receptor for AGEs; **RNS:** Reactive nitrogen species; **RO[•]:** Alkoxy radical; **RO₂^{•-}:** Peroxyl radicals; **ROS:** Reactive oxygen species; **RSS:** Reactive sulphur species; **SCD:** Sickle cell disease; **SDS:** Sodium

dodecyl sulfate; **Se**: Selenium; **SET**: Single electron transfer; **SNpc**: Substantia nigra pars compacta; **SOD**: Superoxide dismutase; **TBARS**: Thiobarbituric acid reactive substance; **TBHQ**: Tertiary butyl hydroquinone; **TCA**: Tricarboxylic acid; **TGF- β** : Transforming growth factor- β ; **TOSCA**: Total oxyradical scavenging capacity assay; **TRAP**: Total radical-trapping antioxidant parameter; **TrxR**: Thioredoxin reductase; **Tyr**: Tyrosine; **UDP-GlcNAc**: UDP-N-acetylglucosamine; **VCAM-1**: Vascular cell adhesion molecule-1; **Zn**: Zinc; **γ -GGT**: γ -Glutamyl transpeptidase; **8-OHdG**: 8-hydroxy-2'-deoxyguanosine.

REFERENCES

- Nimse SB, Pal D. Free radicals, natural antioxidants, and their reaction mechanisms. *RSC Adv.* 2015; 2015:20.
- Athukorala Y, Lee KW, Song C, Ahn CB, Shin TS, Cha YJ, *et al.* Potential antioxidant activity of marine red alga *Grateloupia filicina* extracts. *J Food Lipids.* 2003; 2003:20.
- Shahidi F, Zhong Y. Measurement of antioxidant activity. *J Funct Foods.* 2015; 2015:20.
- Liyana-Pathirana C, Dexter J, Shahidi F. Antioxidant properties of wheat as affected by pearling. *J Agric Food Chem.* 2006; 2006:20.
- Shahidi F, Alasalvar C, Liyana-Pathirana CM. Antioxidant phytochemicals in hazelnut (*Corylus avellana* L.) and its by-products. *J Agric Food Chem.* 2007; 2007:20.
- Cumby N, Zhong Y, Naczek M, Shahid F. Antioxidant activity and water-holding capacity of canola protein hydrolysates. *Food Chem.* 2008; 2008:20.
- Lobo V, Patil A, Phatak A, Chandra N. Free radicals, antioxidants and functional foods: Impact on human health. *Pharmacogn Rev.* 2010; 2010:20.
- Lü JM, Lin PH, Yao Q, Chen C. Chemical and molecular mechanisms of antioxidants: experimental approaches and model systems. *J Cell Mol Med.* 2010; 2010:20.
- Vajragupta O, Boonchoong P, Berliner LJ. Manganese complexes of curcumin analogues: evaluation of hydroxyl radical scavenging ability, superoxide dismutase activity and stability towards hydrolysis. *Free Radic Res.* 2004; 2004:20.
- Giles GI, Jacob C. Reactive sulfur species: an emerging concept in oxidative stress. *Biol Chem.* 2002; 2002:20.
- Geier DA, Kern JK, Garver CR, Adams JB, Audhya T, Geier MR. A prospective study of transsulfuration biomarkers in autistic disorders. *Neurochem Res.* 2009; 2009:20.
- Geier DA, Kern JK, Garver CR, Adams JB, Audhya T, Nataf R, *et al.* Biomarkers of environmental toxicity and susceptibility in autism. *J Neurol Sci.* 2009; 2009:20.
- Prior RL, Wu X, Schaich K. Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *J Agric Food Chem.* 2005; 2005:20.
- Shahidi F, Zhong Y. Novel antioxidants in food quality preservation and health promotion. *Eur J Lipid Sci Technol.* 2010; 2010:20.
- Aziz MA, Diab AS, Mohammed AA. Antioxidant categories and mode of action. *Antioxidants.* 2019. doi:10.5772/intechopen.83544
- Hurrell RF. Influences of vegetable protein sources on trace element and mineral bioavailability. *J Nutr.* 2003; 2003:20.
- Mathew BB, Tiwari A, Jatawa SK. Free radicals and antioxidants: A review. *J Pharm Res.* 2011; 2011:20.
- Adwas AA, Elsayed ASI, Azab AE, Quwaydir FA. Oxidative stress and antioxidant mechanisms in human system. *J Appl Biotechnol Bioeng.* 2019; 2019:20.
- Apak R, Gorinstein S, Böhm V, Schaich Karen M, Özyürek M, Güçlü K. Methods of measurement and evaluation of natural antioxidant capacity/activity (IUPAC Technical Report). *Pure Appl Chem.* 2013; 2013:20.
- Fox SI. *Human Physiology.* 16th ed. New York: McGraw-Hill Education; 2017. p. 666-70.
- Babior BM. NADPH oxidase: an update. *Blood.* 1999; 1999:19.
- Knight JA. Free radicals, antioxidants, aging and disease. Washington: AACCC Press; 1999.
- Zhu L, Gunn C, Beckman JS. Bactericidal activity of peroxynitrite. *Arch Biochem Biophys.* 1992; 1992:19.
- Valko M, Leibfriz D, Moncol J, Cronin MT, Mazur M, Telser J. Free radicals and antioxidants in normal physiological functions and human disease. *Int J Biochem Cell Biol.* 2007; 2007:20.
- Lin HC, Tsai SH, Chen CS, Chang YC, Lee CM, Lai ZY, *et al.* Structure-activity relationship of coumarin derivatives on xanthine oxidase-inhibiting and free radical-scavenging activities. *Biomed Pharmacol.* 2008; 2008:20.
- Schroder K, Vecchione C, Jung O, Schreiber JG, Shiri-Sverdlov R, van Gorp PJ, *et al.* Xanthine oxidase inhibitor tungsten prevents the development of atherosclerosis in ApoE knockout mice fed a Western-type diet. *Free Radic Biol Med.* 2006; 2006:20.
- Krishnamurthy P, Wadhvani A. Antioxidant enzymes and human health. In: El-Missiry MA, editor. *Antioxidant Enzyme.* Croatia: In Tech; 2012. p. 3-18.
- Castilla P, Dávalos A, Teruel JL, Cerrato F, Fernández-Lucas M, Merino JL, *et al.* Comparative effects of dietary supplementation with red grape juice and α -tocopherol on production of superoxide by circulating neutrophil NADPH oxidase in hemodialysis patients. *Am J Clin Nutr.* 2008; 2008:20.
- Voora D, McLeod HL, Eby C, Gage BF. The pharmacogenetics of coumarin therapy. *Pharmacogenomics.* 2005; 2005:20.
- Scandalios JG. Oxidative stress: molecular perception and transduction of signals triggering antioxidant gene defenses. *Braz J Med Biol Res.* 2005; 2005:20.
- Pandey KB, Rizvi SI. Markers of oxidative stress in erythrocytes and plasma during aging in humans. *Oxid Med Cell Longev.* 2010; 2010:20.
- Turrens JF, Crapo JD, Freeman BA. Protection against oxygen toxicity by intravenous injection of liposome-entrapped catalase and superoxide dismutase. *J Clin Invest.* 1984; 1984:19.
- Li Y, Huang TT, Carlson EJ, Melov S, Ursell PC, Olson JL, *et al.* Dilated cardiomyopathy and neonatal lethality in mutant mice lacking manganese superoxide dismutase. *Nat Genet.* 1995; 1995:19.
- Van Loon APGM, Pesold-Hurt B, Schatz G. A yeast mutant lacking mitochondrial manganese-superoxide dismutase is hypersensitive to oxygen. *Proc Natl Acad Sci USA.* 1986; 1986:19.
- Vitale M, Di Matola T, D'Ascoli F. Iodide excess induces apoptosis in thyroid cells through a p53-independent mechanism involving oxidative stress. *Endocrinol Soc.* 2000; 2000:20.
- James SJ, Rose S, Melnyk S, Jernigan S, Blossom S, Pavliv O, *et al.* Cellular and mitochondrial glutathione redox imbalance in lymphoblastoid cells derived from children with autism. *FASEB J.* 2009; 2009:20.
- Topal F, Nar M, Gocer H, Kalin P, Kocyigit UM, Gülçin İ, *et al.* Antioxidant activity of taxifolin: An activity-structure relationship. *J Enzyme Inhib Med Chem.* 2016; 2016:20.
- Ak T, Gulcin I. Antioxidant and radical scavenging properties of curcumin. *Chem Biol Interact.* 2008; 2008:20.
- Kazazica SP, Butkovic V, Srazica D, Klasinc L. Gas-phase ligation of Fe⁺ and Cu⁺ ions with some flavonoids. *J Agric Food Chem.* 2006; 2006:20.
- Cakmak CK, Gulcin I. Anticholinergic and antioxidant activities of usnic acid-An activity-structure insight. *Toxicol Rep.* 2019; 2019:20.
- Gulcin I. Antioxidant properties of resveratrol: A structure-activity insight. *Innov Food Sci Emerg Technol.* 2010; 2010:20.
- Fiorucci SB, Golebiowski J, Cabrol-Bass D, Antonczak S. DFT study of quercetin activated forms involved in antiradical, antioxidant, and prooxidant biological processes. *J Agric Food Chem.* 2007; 2007:20.
- Cherrak SA, Mokhtari-Soulimane N, Berroukeche F, Bensenane B, Cherbonnel A, Merzouk H, *et al.* *In vitro* Antioxidant versus Metal Ion Chelating Properties of Flavonoids: A Structure-Activity Investigation. *PLoS ONE.* 2016; 11(10): e0165575.
- Gulcin I. Antioxidant activity of caffeic acid (3,4-dihydroxycinnamic acid). *Toxicology.* 2006; 2006:20.
- Gocer H, Gulcin I. Caffeic acid phenethyl ester (CAPE): Correlation of structure and antioxidant properties. *Int J Food Sci.* 2011; 2011:20.
- Lindsay D, Kerr W. Cobalt close-up. *Nat Chem.* 2014; 2014:20.
- Eruygur N, Atas M, Tekin M, Taslimi P, Kocyigit UM, Gulcin I. *In vitro* antioxidant, antimicrobial, anticholinesterase and antidiabetic activities of Turkish endemic *Achillea cucullata* (Asteraceae) from ethanol extract. *S Afr J Bot.* 2019; 2019:20.
- Ghosh N, Chakraborty T, Mallick S, Mana S, Singha D, Ghosh B, *et al.* Synthesis, characterization and study of antioxidant activity of quercetin-magnesium complex. *Spectrochim Acta A Mol Biomol Spectrosc.* 2015; 2015:20.
- Han H, Yilmaz H, Gulcin I. Antioxidant activity of flaxseed (*Linum usitatissimum* L.) and analysis of its polyphenol contents by LC-MS/MS. *Rec Nat Prod.* 2018; 2018:20.
- Kalin P, Gulcin I, Goren AC. Antioxidant activity and polyphenol content of cranberries (*Vaccinium macrocarpon*). *Rec Nat Prod.* 2015; 2015:20.
- Gulcin I, Alwasel SH. Metal Ions, Metal Chelators and Metal Chelating Assay as Antioxidant Method. *Processes.* 2022; 10.
- Pacher P, Beckman JS, Liaudet L. Nitric oxide and peroxynitrite in health and disease. *Physiol Rev.* 2007; 2007:20.
- Beckman JS, Koppenol WH. Nitric oxide, superoxide and peroxynitrite: the good, the bad and ugly. *Am J Physiol.* 1996; 1996:19.
- Eiserich JP, Cross CE, Jones AD, Halliwell B, van der Vliet A. Formation of nitrating and chlorinating species by reaction of nitrite with hypochlorous acid. A novel mechanism for nitric oxide-mediated protein modification. *J Biol Chem.* 1996; 1996:19.
- Mohiuddin I, Chai H, Lin PH, Lumsden AB, Yao Q, Chen C. Nitrotyrosine and chlorotyrosine: clinical significance and biological functions in the vascular system. *J Surg Res.* 2006; 2006:20.
- Viappiani S, Schulz R. Detection of specific nitrotyrosine-modified proteins as a marker of oxidative stress in cardiovascular disease. *Am J Physiol Heart Circ Physiol.* 2006; 290:H2167-8.
- Alkam T, Nitta A, Mizoguchi H, Itoh A, Nabeshima T. A natural scavenger of peroxynitrites, rosmarinic acid, protects against impairment of memory induced by Abeta. *Behav Brain Res.* 2007; 2007:20.
- Medina-Campos ON, Barrera D, Segoviano-Murillo S, Rocha D, Maldonado PD, Mendoza-Patiño N, *et al.* S-allylcysteine scavenges singlet oxygen and hypochlorous

- acid and protects LLC-PK(1) cells of potassium dichromate-induced toxicity. *Food Chem Toxicol.* 2007; 2007:20.
59. Fernandes E, Toste SA, Lima JL, Reis S. The metabolism of sulindac enhances its scavenging activity against reactive oxygen and nitrogen species. *Free Radic Biol Med.* 2003; 2003:20.
 60. Zheng LF, Wei QY, Cai YJ, Fang JG, Zhou B, Yang L, *et al.* DNA damage induced by resveratrol and its synthetic analogues in the presence of Cu (II) ions: mechanism and structure-activity relationship. *Free Radic Biol Med.* 2006; 2006:20.
 61. Ramoutar RR, Brumaghim JL. Effects of inorganic selenium compounds on oxidative DNA damage. *J Inorg Biochem.* 2007; 2007:20.
 62. Zhao F, Tang YZ, Liu ZQ. Protective effect of icariin on DNA against radical-induced oxidative damage. *J Pharm Pharmacol.* 2007; 2007:20.
 63. Kang HS, Kim KR, Jun EM, Park SH, Lee TS, Suh JW, *et al.* Cyathuscavins A, B, and C, new free radical scavengers with DNA protection activity from the *Basidiomycete Cyathus stercoreus*. *Bioorg Med Chem Lett.* 2008; 2008:20.
 64. Zhao F, Liu ZQ, Wu D. Antioxidative effect of melatonin on DNA and erythrocytes against free-radical-induced oxidation. *Chem Phys Lipids.* 2008; 2008:20.
 65. Perron NR, Hodges JN, Jenkins M, Brumaghim JL. Predicting how polyphenol antioxidants prevent DNA damage by binding to iron. *Inorg Chem.* 2008; 2008:20.
 66. Kang KS, Kim HY, Pyo JS, Yokozawa T. Increase in the free radical scavenging activity of ginseng by heat-processing. *Biol Pharm Bull.* 2006; 2006:20.
 67. Dai F, Chen WF, Zhou B. Antioxidant synergism of green tea polyphenols with alpha-tocopherol and L-ascorbic acid in SDS micelles. *Biochimie.* 2008; 2008:20.
 68. Gal S, Lichtenberg D, Bor A, Pinchuk I. Copper-induced peroxidation of phosphatidylserine-containing liposomes is inhibited by nanomolar concentrations of specific antioxidants. *Chem Phys Lipids.* 2007; 2007:20.
 69. Santos-Sánchez NF, Salas-Coronado R, Villanueva-Cañongo C, Hernández-Carlos B. Antioxidants and Their Antioxidant Mechanism. *IntechOpen*; 2019. p. 1-28.
 70. Arouma OI. Assessment of potential prooxidant and antioxidant actions. *J Am Oil Chem Soc.* 1996; 1996:19.
 71. Huang D, Ou B, Prior RL. The chemistry behind antioxidant capacity assays. *J Agric Food Chem.* 2005; 2005:20.
 72. Fukumoto LR, Mazza G. Assessing antioxidant and pro-oxidant activities of phenolic compounds. *J Agric Food Chem.* 2000; 2000:20.
 73. Lichtenthaler R, Marx F. Total oxidant scavenging capacities of common European fruit and vegetable juices. *J Agric Food Chem.* 2005; 2005:20.
 74. Benzie IFF, Strain JJ. Ferric reducing/antioxidant power assay: Direct measure of total antioxidant activity of biological fluids and modified version for simultaneous measurement of total antioxidant power and ascorbic acid concentration. *Methods Enzymol.* 1999; 1999:19.
 75. Kedare SB, Singh RP. Genesis and development of DPPH method of antioxidant assay. *J Food Sci Technol.* 2011; 2011:20.
 76. Dawidowicz AL, Wianowska D, Olszowy M. On practical problems in estimation of antioxidant activity of compounds by DPPH method (problems in estimation of antioxidant activity). *Food Chem.* 2012; 2012:20.
 77. Musa KH, Abdullah A, Kuswandi B, Hidayat MA. A novel high throughput method based on the DPPH dry reagent array for determination of antioxidant activity. *Food Chem.* 2013; 2013:20.
 78. Gardner PT, McPhaila DB, Duthie GG. Electron spin resonance spectroscopic assessment of the antioxidant potential of teas in aqueous and organic media. *J Sci Food Agric.* 1998; 1998:19.
 79. Zhong Y, Shahidi F. Lipophilized epigallocatechin gallate (EGCG) derivatives as novel antioxidants. *J Agric Food Chem.* 2011; 2011:20.
 80. Amatongchai M, Laosing S, Chailapakul O, Nacapricha D. Simple flow injection for screening of total antioxidant capacity by amperometric detection of DPPH radical on carbon nanotube modified-glassy carbon electrode. *Talanta.* 2012; 2012:20.
 81. Sireerat I, Schulte A. Automated electrochemical free radical scavenger screening in dietary samples. *Anal Chem.* 2012; 2012:20.
 82. Manzocco L, Anese M, Nicoli MC. Antioxidant properties of tea extracts as affected by processing. *Lebensm Wiss Technol.* 1998; 1998:19.
 83. Ghiselli A, Serafini M, Maiani G, Azzini E, Ferro-Luzzi A. A fluorescence-based method for measuring total plasma antioxidant capability. *Free Radic Biol Med.* 1995; 1995:19.
 84. Bartosz G, Janaszewska A, Ertel D, Bartosz M. Simple determination of peroxy radical-trapping capacity. *Int Union Biochem Mol Biol.* 1998; 1998:19.
 85. Valkonen M, Kuusi T. Spectrophotometric assay for total peroxy radical-trapping antioxidant potential in human serum. *J Lipid Res.* 1997; 1997:19.
 86. Santos-Sánchez NF, Salas-Coronado R, Villanueva-Cañongo C, Hernández-Carlos B. Antioxidant Compounds and Their Antioxidant Mechanism. *Antioxidants.* 2019.
 87. Cao G, Sofic E, Prior RL. Antioxidant and pro-oxidant behavior of flavonoids: Structure-activity relationships. *Free Radic Biol Med.* 1997; 1997:19.
 88. Seeram NP, Aviram M, Zhang Y, Henning SM, Feng L, Dreher M, *et al.* Comparison of antioxidant capacity of commonly consumed polyphenol-rich beverages in the United States. *J Agric Food Chem.* 2008; 2008:20.
 89. Jacobo-Velázquez DA, Cisneros-Zeballos L. Correlations of antioxidant activity against phenolic content revisited: A new approach in data analysis for food and medicinal plants. *J Food Sci.* 2009; 2009:20.
 90. Huang D, Ou B, Hampsch-Woodill M, Flanagan J, Prior R. High-throughput assay of oxygen radical absorbance capacity (ORAC) using a multichannel liquid handling system coupled with a microplate fluorescence reader in 96-well format. *J Agric Food Chem.* 2002; 2002:20.
 91. Prior RL, Hoang H, Gu L. Assays for hydrophilic and lipophilic antioxidant capacity (oxygen radical absorbance capacity (ORAC (FL)) of plasma and other biological and food samples. *J Agric Food Chem.* 2003; 2003:20.
 92. Alam MN, Bristi NJ, Rafiquzzaman M. Review on *in vivo* and *in vitro* methods evaluation of antioxidant activity. *Saudi Pharm J.* 2013; 2013:20.
 93. Ohkawa H, Onishi N, Yagi K. Assay for lipid peroxidation in animal tissue by thiobarbituric acid reaction. *Anal Biochem.* 1979; 1979:19.
 94. Zimmermann AM, Rivero AC, Bezerra AS, Ruviaro AR, Novack MME, Nörnberg JL. Antioxidant potential of barley extract in rats subjected to a high-fat diet. *Food Sci Technol Campinas.* 2013; 2013:20.
 95. Chikezie PC, Uwakwe AA. Protective effect of *Allium sativa* extract against carbon tetrachloride-induced hepatic oxidative stress and hyperlipidemia in rats. *Afr J Biotechnol.* 2014; 2014:20.
 96. Benzie IFF, Strain JJ. The ferric reducing ability of plasma (FRAP) as a measure of 'antioxidant power': The FRAP assay. *Anal Biochem.* 1996; 1996:19.
 97. Sapakal VD, Shikalgar TS, Ghadge RV, Adnalk RS, Naikwade NS, Magdum CS. *In vivo* screening of antioxidant profile: a review. *J Herb Med Toxicol.* 2008; 2008:20.
 98. Ellman GL. Tissue sulfhydryl groups. *Arch Biochem Biophys.* 1959; 1959:19.
 99. Wood JL. In: Fishman WH, editor. *Metabolic Conjugation and Metabolic Hydrolysis.* Vol. 2. New York: Academic Press; 1970. p. 61-299.
 100. Jocelyn PC. *Biochemistry of the SH Group.* London: Academic Press; 1972. p. 10.
 101. Habig WH, Pabst MJ, Jacoby WB. Glutathione S-transferases: the first enzymatic step in mercapturic acid formation. *J Biol Chem.* 1974; 1974:19.
 102. McCord J, Fridovich I. Superoxide dismutase, an enzymic function for erythrocytes. *J Biol Chem.* 1969; 1969:19.
 103. Aebi H. Catalase. *Methods Enzymol.* 1984; 1984:19.
 104. Singhal PK, Kapur N, Dhillon KS, Beamish RE, Dhalla NS. Role of free radicals in catecholamine cardiomyopathy. *Can J Physiol Pharmacol.* 1982; 1982:19.
 105. Kakkar P, Das B, Viswanathan PN. A modified spectrophotometric assay of superoxide dismutase. *Indian J Biochem Biophys.* 1984; 1984:19.
 106. Wohaieb SA, Godin DV. Alterations in free radical tissue-defense mechanisms in streptozotocin-induced diabetes in rat: Effects of insulin treatment. *Diabetes.* 1987; 1987:19.
 107. Yakubu MA, Sofola OA, Igbo I, Oyekan AO. Link between free radicals and protein kinase C in glucose-induced alteration of vascular dilation. *Life Sci.* 2004; 2004:20.
 108. Shaikh H, Shrivastava VK. Effects of streptozotocin induced diabetes mellitus type 1 on the rat brain antioxidant status and activity of acetylcholinesterase: a novel and potential treatment by *Vitex negundo*. *Int J Pharm Pharm Sci.* 2014; 2014:20.
 109. Styskal JL, Van Remmen H, Richardson A, Salmon AB. Oxidative stress and diabetes: what can we learn about insulin resistance from antioxidant mutant mouse models? *Free Radic Biol Med.* 2012; 2012:20.
 110. Chikezie PC, Ojiako OA, Ogbuji AC. Oxidative stress in diabetes mellitus. *Int J Biol Chem.* 2015; 2015:20.
 111. Pickering RJ, Rosado CJ, Sharma A, Buksh S, Tate M, de Haan JB. Recent novel approaches to limit oxidative stress and inflammation in diabetic complications. *Clin Transl Immunol.* 2018; 7:e1016.
 112. Alfadda AA, Sallam RM. Reactive oxygen species in health and disease. *J Biomed Biotechnol.* 2012; 2012:20 pages.
 113. Taverne YJ, Bogers AJ, Duncker DJ, Merkus D. Reactive oxygen species and the cardiovascular system. *Oxid Med Cell Longev.* 2013; 2013:20.
 114. Sharif-Rad M, Anil Kumar NV, Zucca P, *et al.* Lifestyle, oxidative stress, and antioxidants: back and forth in the pathophysiology of chronic diseases. *Front Physiol.* 2020; 2020:20.
 115. Elahi MM, Kong YX, Matata BM. Oxidative stress as a mediator of cardiovascular disease. *Oxid Med Cell Longev.* 2009; 2009:20.
 116. Singh RB, Mengi SA, Xu YJ, Arneja AS, Dhalla NS. Pathogenesis of atherosclerosis: a multifactorial process. *Exp Clin Cardiol.* 2002; 2002:20.
 117. Perrotta I, Aquila S. The role of oxidative stress and autophagy in atherosclerosis. *Oxid Med Cell Longev.* 2015; 2015:20.
 118. Esper RJ, Nordaby RA, Vilarino JO, Paragano A, Cacharron JL, Machado RA. Endothelial dysfunction: a comprehensive appraisal. *Cardiovasc Diabetol.* 2006; 2006:20.
 119. Förstermann U. Nitric oxide and oxidative stress in vascular disease. *Pflügers Arch-Eur J Physiol.* 2010; 2010:20.
 120. Montezano AC, Touyz RM. Reactive oxygen species and endothelial function-role of nitric oxide synthase uncoupling and Nox family nicotinamide adenine dinucleotide phosphate oxidases. *Basic Clin Pharmacol Toxicol.* 2012; 2012:20.
 121. Kibel A, Lukinac AM, Dambic V, Juric I, Selthofer-Relatic K. Oxidative stress in ischemic heart disease. *Oxid Med Cell Longev.* 2020; 2020:20.
 122. Li H, Horke S, Förstermann U. Vascular oxidative stress, nitric oxide and atherosclerosis. *Atherosclerosis.* 2014; 2014:20.
 123. Higashi Y, Sasaki S, Nakagawa K, Matsuura H, Oshima T, Chayama K. Endothelial function and oxidative stress in renovascular hypertension. *N Engl J Med.* 2002; 2002:20.

124. Lip GYH, Edmunds E, Nuttall SL, Landray MJ, Blann AD, Beevers DG. Oxidative stress in malignant and non-malignant phase hypertension. *J Hum Hypertens.* 2002; 2002:20.
125. He F, Zuo L. Redox roles of reactive oxygen species in cardiovascular diseases. *Int J Mol Sci.* 2015; 2015:20.
126. Guzik TJ, West NE, Black E, *et al.* Vascular superoxide production by NAD(P)H oxidase: Association with endothelial dysfunction and clinical risk factors. *Circ Res.* 2000; 2000:20.
127. Azarsiz E, Kayikcioglu M, Payzin S, Yildirim Sözmen E. PON1 activities and oxidative markers of LDL in patients with angiographically proven coronary artery disease. *Int J Cardiol.* 2003; 2003:20.
128. Hare JM, Stamler JS. NO/redox disequilibrium in the failing heart and cardiovascular system. *J Clin Invest.* 2005; 2005:20.
129. Wattanapitayakul S K, Bauer JA. Oxidative pathways in cardiovascular disease: roles, mechanisms, and therapeutic implications. *Pharmacol Ther.* 2001; 2001:20.
130. Gutteridge JM, Halliwell B. Free radicals and antioxidants in the year 2000: A historical look to the future. *Ann NY Acad Sci.* 2000; 2000:20.
131. Matough FA, Budin SB, Hamid ZA, Alwahaibi N, Mohamed J. The role of oxidative stress and antioxidants in diabetic complications. *Sultan Qaboos Univ Med J.* 2012; 2012:20.
132. Wolff SP, Jiang ZY, Hunt JV. Protein glycation and oxidative stress in diabetes mellitus and ageing. *Free Radic Biol Med.* 1991; 1991:19.
133. Ceriello A, Quatraro A, Giugliano D. New insights on non-enzymatic glycosylation may lead to therapeutic approaches for the prevention of diabetic complications. *Diabet Med.* 1992; 1992:19.
134. Vaag A, Damsbo P, Hother-Nielsen O, Beck-Nielsen H. Hyperglycaemia compensates for the defects in insulin-mediated glucose metabolism and in the activation of glycogen synthase in the skeletal muscle of patients with type 2 (non-insulin-dependent) diabetes mellitus. *Diabetologia.* 1992; 1992:19.
135. Giacco F, Brownlee M. Oxidative stress and diabetic complications. *Circ Res.* 2010; 2010:20.
136. Di Naso FC, Dias AS, Porawski M, Marroni NAP. Exogenous superoxide dismutase: action on liver oxidative stress in animals with streptozotocin-induced diabetes. *Exp Diabetes Res.* 2011; 2011:20.
137. Aronson D, Rayfield EJ. How hyperglycemia promotes atherosclerosis: Molecular mechanisms. *Cardiovasc Diabetol.* 2002; 2002:20.
138. Wallace DC. Diseases of the mitochondrial DNA. *Annu Rev Biochem.* 1992; 1992:19.
139. Brownlee M. Biochemistry and molecular cell biology of diabetic complications. *Nature.* 2001; 2001:20.
140. Brownlee M. The pathobiology of diabetic complications: A unifying mechanism. *Diabetes.* 2005; 2005:20.
141. Papachristoforou E, Lambadiari V, Maratou E, Makrilakis K. Association of glycemic indices (hyperglycemia, glucose variability, and hypoglycemia) with oxidative stress and diabetic complications. *J Diabetes Res.* 2020; 2020:20.
142. Du X, Matsumura T, Edelstein D, *et al.* Inhibition of GAPDH activity by poly(ADP-ribose) polymerase activates three major pathways of hyperglycemic damage in endothelial cells. *J Clin Invest.* 2003; 2003:20.
143. Mikhed Y, Daiber A, Steven S. Mitochondrial oxidative stress, mitochondrial DNA damage and their role in age-related vascular dysfunction. *Int J Mol Sci.* 2015; 2015:20.
144. Zakaria EM, El-Bassossy HM, El-Maraghy NN, Ahmed AF, Ali AA. PARP-1 inhibition alleviates diabetic cardiac complications in experimental animals. *Eur J Pharmacol.* 2016; 2016:20.
145. Qin WD, Liu GL, Wang J, *et al.* Poly(ADP-ribose) polymerase 1 inhibition protects cardiomyocytes from inflammation and apoptosis in diabetic cardiomyopathy. *Oncotarget.* 2016; 2016:20.
146. Sultan CS, Saackel A, Stank A, *et al.* Impact of carbonylation on glutathione peroxidase-1 activity in human hyperglycemic endothelial cells. *Redox Biol.* 2018; 2018:20.
147. Davison GW, George L, Jackson SK, *et al.* Exercise, free radicals, and lipid peroxidation in type 1 diabetes mellitus. *Free Radic Biol Med.* 2002; 2002:20.
148. Ramakrishna V, Jailkhani R. Evaluation of oxidative stress in insulin dependent diabetes mellitus (IDDM) patients. *Diagn Pathol.* 2007; 2007:20.
149. Godin DV, Wohaiab SA, Garnett ME, Goumeniouk AD. Antioxidant enzyme alterations in experimental and clinical diabetes. *Mol Cell Biochem.* 1988; 1988:19.
150. Ohkuwa T, Sato Y, Naoi M. Hydroxyl radical formation in diabetic rats induced by streptozotocin. *Life Sci.* 1995; 1995:19.
151. Ceolotto G, Bevilacqua M, Papparella I, *et al.* Insulin generates free radicals by an NAD(P)H, phosphatidylinositol-3'-kinase-dependent mechanism in human skin fibroblasts *ex vivo*. *Diabetes.* 2004; 2004:20.
152. Klaunig JE, Kamendulis LM, Hocevar BA. Oxidative stress and oxidative damage in carcinogenesis. *Toxicol Pathol.* 2010; 2010:20.
153. Glasauer A, Chandel NS. Targeting antioxidants for cancer therapy. *Biochem Pharmacol.* 2014; 2014:20.
154. Pizzino G, Bitto A, Interdonato M, *et al.* Oxidative stress and DNA repair and detoxification gene expression in adolescents exposed to heavy metals living in the Milazzo-Valle del Mela area (Sicily, Italy). *Redox Biol.* 2014; 2014:20.
155. Pizzino G, Irrera N, Cucinotta M, *et al.* Oxidative stress: Harms and benefits for human health. *Oxid Med Cell Longev.* 2017; 2017:20.
156. Liou GY, Döppler H, DelGiorno KE, *et al.* Mutant KRas-induced mitochondrial oxidative stress in acinar cells upregulates EGFR signaling to drive formation of pancreatic precancerous lesions. *Cell Rep.* 2016; 2016:20.
157. Herrera B, Murillo MM, Alvarez-Barrientos A, Beltran J, Fernandez M, Fabregat I. Source of early reactive oxygen species in the apoptosis induced by transforming growth factor-beta in fetal rat hepatocytes. *Free Radic Biol Med.* 2004; 2004:20.
158. Cortat B, Garcia CCM, Quinet A, Schuch AP, De Lima-Bessa KM, Menck CFM. The relative roles of DNA damage induced by UVA irradiation in human cells. *Photochem Photobiol Sci.* 2013; 2013:20.
159. Saha SK, Lee SB, Won J, *et al.* Correlation between oxidative stress, nutrition, and cancer initiation. *Int J Mol Sci.* 2017; 2017:20.
160. Sander CS, Hamm F, Elsner P, Thiele JJ. Oxidative stress in malignant melanoma and non-melanoma skin cancer. *Br J Dermatol.* 2003; 2003:20.
161. Sova H, Jukkola-Vuorinen A, Puistola U, Kauppila S, Karihtala P. 8-Hydroxydeoxyguanosine: a new potential independent prognostic factor in breast cancer. *Br J Cancer.* 2010; 2010:20.
162. Forcados GE, James DB, Sallau AB, Muhammad A, Mabeta P. Oxidative stress and carcinogenesis: potential of phytochemicals in breast cancer therapy. *Nutr Cancer.* 2017; 2017:20.
163. Miyake H, Hara I, Kamidono S, Eto H. Oxidative DNA damage in patients with prostate cancer and its response to treatment. *J Urol.* 2004; 2004:20.
164. Weiss JM, Goode EL, Ladiges WC, Ulrich CM. Polymorphic variation in hOGG1 and risk of cancer: A review of the functional and epidemiologic literature. *Mol Carcinog.* 2005; 2005:20.
165. Diakowska D, Lewandowski A, Kopec W, Diakowski W, Chrzanoska T. Oxidative DNA damage and total antioxidant status in serum of patients with esophageal squamous cell carcinoma. *Hepatogastroenterol.* 2007; 2007:20.
166. Tanaka H, Fujita N, Sugimoto R, *et al.* Hepatic oxidative DNA damage is associated with increased risk for hepatocellular carcinoma in chronic hepatitis C. *Br J Cancer.* 2008; 2008:20.
167. Gottschling BC, Maronpot RR, Hailey JR, *et al.* The role of oxidative stress in indium phosphide-induced lung carcinogenesis in rats. *Toxicol Sci.* 2001; 2001:20.
168. Muguruma M, Unami A, Kanki M, *et al.* Possible involvement of oxidative stress in piperonyl butoxide induced hepatocarcinogenesis in rats. *Toxicology.* 2007; 2007:20.
169. Jaramillo MC, Zhang DD. The emerging role of the Nrf2-Keap1 signaling pathway in cancer. *Genes Dev.* 2013; 2013:20.
170. Aikawa R, Komuro I, Yamazaki T, *et al.* Oxidative stress activates extracellular signal-regulated kinases through Src and Ras in cultured cardiac myocytes of neonatal rats. *J Clin Invest.* 1997; 1997:19.
171. Querfurth HW, LaFerla FM. Alzheimer's disease. *N Engl J Med.* 2010; 2010:20.
172. Goedert M, Spillantini MG. A century of Alzheimer's disease. *Science.* 2006; 2006:20.
173. Rak M, Bénit P, Chrétien D, *et al.* Mitochondrial cytochrome c oxidase deficiency. *Clin Sci.* 2016; 2016:20.
174. Zhang XD, Wang Y, Wu JC, *et al.* Down-regulation of Bcl-2 enhances autophagy activation and cell death induced by mitochondrial dysfunction in rat striatum. *J Neurosci Res.* 2009; 2009:20.
175. Elgenaidi IS, Spiers JP. Regulation of the phosphoprotein phosphatase 2A system and its modulation during oxidative stress: A potential therapeutic target? *Pharmacol Ther.* 2019; 2019:20.
176. Toral-Rios D, Pichardo-Rojas PS, Alonso-Vanegas M, Campos-Pena V. GSK3beta and tau protein in Alzheimer's disease and epilepsy. *Front Cell Neurosci.* 2020; 2020:20.
177. Misrani A, Tabassum S, Yang L. Mitochondrial dysfunction and oxidative stress in Alzheimer's disease. *Front Aging Neurosci.* 2021; 2021:20.
178. Hirai K, Aliev G, Nunomura A, *et al.* Mitochondrial abnormalities in Alzheimer's disease. *J Neurosci.* 2001; 2001:20.
179. Zhu X, Perry G, Moreira PI, *et al.* Mitochondrial abnormalities and oxidative imbalance in Alzheimer disease. *J Alzheimers Dis.* 2006; 2006:20.
180. Mutisya EM, Bowling AC, Beal MF. Cortical cytochrome oxidase activity is reduced in Alzheimer's disease. *J Neurochem.* 1994; 1994:19.
181. Kumar V, Khan AA, Tripathi A, Dixit PK, Bajaj U. Role of oxidative stress in various diseases: Relevance of dietary antioxidants. *J Phytopharmacol.* 2015; 2015:20.
182. Matsuoka Y, Picciano M, La Franco J, Duff K. Fibrillar β -amyloid evokes oxidative damage in a transgenic mouse model of Alzheimer's disease. *Neuroscience.* 2001; 2001:20.
183. Apelt J, Bigl M, Wunderlich P, Schliebs R. Aging-related increase in oxidative stress correlates with developmental pattern of beta-secretase activity and beta-amyloid plaque formation in transgenic Tg2576 mice with Alzheimer-like pathology. *Int J Dev Neurosci.* 2004; 2004:20.
184. Manczak M, Anekonda TS, Henson E, Park BS, Quinn J, Reddy PH. Mitochondria are a direct site of $A\beta$ accumulation in Alzheimer's disease neurons: implications for free radical generation and oxidative damage in disease progression. *Hum Mol Genet.* 2006; 2006:20.
185. Mohammad-Abdul H, Sultana R, Keller JN, St. Clair DK, Markesbery WR, Butterfield DA. Mutations in amyloid precursor protein and presenilin-1 genes increase the basal oxidative stress in murine neuronal cells and lead to increased sensitivity to oxidative

- stress mediated by amyloid β -peptide (1-42), H₂O₂ and kainic acid: implications for Alzheimer's disease. *J Neurochem*. 2006; 2006:20.
186. De Felice FG, Velasco PT, Lambert MP, *et al.* Abeta oligomers induce neuronal oxidative stress through an N-methyl-D-aspartate receptor-dependent mechanism that is blocked by the Alzheimer drug memantine. *J Biol Chem*. 2007; 2007:20.
 187. Deibel MA, Ehmann WD, Markesbery WR. Copper, iron, and zinc imbalances in severely degenerated brain regions in Alzheimer's disease: possible relation to oxidative stress. *J Neurol Sci*. 1996; 1996:19.
 188. Kenche VB, Barnham KJ. Alzheimer's disease & metals: therapeutic opportunities. *Br J Pharmacol*. 2011; 2011:20.
 189. Lee JY, Mook-Jung I, Koh JY. Histochemically reactive zinc in plaques of the Swedish mutant beta-amyloid precursor protein transgenic mice. *J Neurosci*. 1999; 19(11):RC10.
 190. Zhang J, Liu Q, Chen Q, *et al.* Nicotine attenuates beta-amyloid-induced neurotoxicity by regulating metal homeostasis. *FASEB J*. 2006; 2006:20.
 191. Dalle-Donne I, Scaloni-Giustarini A, Cavarra D, Tell E, Lungarella G. Proteins as biomarkers of oxidative stress in diseases: the contribution of redox proteomics. *Mass Spectrom Rev*. 2005; 2005:20.
 192. Blesa J, Trigo-Damas I, Quiroga-Varela A, Jackson-Lewis VR. Oxidative stress and Parkinson's disease. *Front Neuroanat*. 2015.
 193. Schapira AHV. Mitochondria in the aetiology and pathogenesis of Parkinson's disease. *Lancet Neurol*. 2008; 2008:20.
 194. Hall CN, Klein-Flügge MC, Howarth C, Attwell D. Oxidative phosphorylation, not glycolysis, powers presynaptic and postsynaptic mechanisms underlying brain information processing. *J Neurosci*. 2012; 2012:20.
 195. Schapira AH, Cooper JM, Dexter D, Clark JB, Jenner P, Marsden CD. Mitochondrial complex I deficiency in Parkinson's disease. *J Neurochem*. 1990; 1990:19.
 196. Hattori N, Tanaka M, Ozawa T, Mizuno Y. Immunohistochemical studies on complexes I, II, III, and IV of mitochondria in Parkinson's disease. *Ann Neurol*. 1991; 1991:19.
 197. Hattingen E, Magerkurth J, Pilatus U, *et al.* Phosphorus and proton magnetic resonance spectroscopy demonstrates mitochondrial dysfunction in early and advanced Parkinson's disease. *Brain*. 2009; 2009:20.
 198. Blin O, Desnuelle C, Rascol O, *et al.* Mitochondrial respiratory failure in skeletal muscle from patients with Parkinson's disease and multiple system atrophy. *J Neurol Sci*. 1994; 1994:19.
 199. Mytilineou C, Werner P, Molinari S, Di Rocco A, Cohen G, Yahr MD. Impaired oxidative decarboxylation of pyruvate in fibroblasts from patients with Parkinson's disease. *J Neural Transm Park Dis Dement Sect*. 1994; 1994:19.
 200. Blandini F, Nappi G, Greenamyre JT. Quantitative study of mitochondrial complex I in platelets of parkinsonian patients. *Mov Disord*. 1998; 1998:19.
 201. Haas RH, Nasirian F, Nakano K, *et al.* Low platelet mitochondrial complex I and complex II/III activity in early untreated Parkinson's disease. *Ann Neurol*. 1995; 1995:19.
 202. Parker WD, Parks JK, Swerdlow RH. Complex I deficiency in Parkinson's disease frontal cortex. *Brain Res*. 2008; 2008:20.
 203. Norris KL, Hao R, Chen LF, *et al.* Convergence of Parkin, PINK1, and α -Synuclein on stress-induced mitochondrial morphological remodeling. *J Biol Chem*. 2015; 2015:20.
 204. van der Merwe C, Jalali Sefid Dashti Z, Christoffels A, Loos B, Bardien S. Evidence for a common biological pathway linking three Parkinson's disease-causing genes: parkin, PINK1 and DJ-1. *Eur J Neurosci*. 2015; 2015:20.
 205. Zuo L, Motherwell MS. The impact of reactive oxygen species and genetic mitochondrial mutations in Parkinson's disease. *Gene*. 2013; 2013:20.
 206. Wang HL, Chou AH, Wu AS, *et al.* PARK6 PINK1 mutants are defective in maintaining mitochondrial membrane potential and inhibiting ROS formation of substantia nigra dopaminergic neurons. *Biochim Biophys Acta*. 2011; 2011:20.
 207. Clark IE, Dodson MW, Jiang C, *et al.* Drosophila pink1 is required for mitochondrial function and interacts genetically with parkin. *Nature*. 2006; 2006:20.
 208. Kitada T, Pisani A, Porter DR, *et al.* Impaired dopamine release and synaptic plasticity in the striatum of PINK1-deficient mice. *Proc Natl Acad Sci USA*. 2007; 2007:20.
 209. Gautier CA, Kitada T, Shen J. Loss of PINK1 causes mitochondrial functional defects and increased sensitivity to oxidative stress. *Proc Natl Acad Sci USA*. 2008; 2008:20.
 210. Goedert M, Spillantini MG, Del Tredici K, Braak H. 100 years of Lewy pathology. *Nat Rev Neurol*. 2013; 2013:20.
 211. Martin LJ, Pan Y, Price AC, *et al.* Parkinson's disease alpha-synuclein transgenic mice develop neuronal mitochondrial degeneration and cell death. *J Neurosci*. 2006; 2006:20.
 212. Dryanovski DI, Guzman JN, Xie Z, *et al.* Calcium entry and α -synuclein inclusions elevate dendritic mitochondrial oxidant stress in dopaminergic neurons. *J Neurosci*. 2013; 2013:20.
 213. Xiang W, Schlachetzki JC, Helling S, *et al.* Oxidative stress-induced posttranslational modifications of alpha-synuclein: specific modification of alpha-synuclein by 4-hydroxy-2-nonenal increases dopaminergic toxicity. *Mol Cell Neurosci*. 2013; 2013:20.
 214. Pukass K, Richter-Landsberg C. Oxidative stress promotes uptake, accumulation, and oligomerization of extracellular α -synuclein in oligodendrocytes. *J Mol Neurosci*. 2014; 2014:20.
 215. Ortuño-Sahagún D, Pallás M, Rojas-Mayorquín AE. Oxidative stress in aging: Advances in proteomic approaches. *Oxid Med Cell Longev*. 2014; 2014:20.
 216. Tsoukalas D, Fragkiadaki P, Docea AO, *et al.* Discovery of potent telomerase activators: Unfolding new therapeutic and anti-aging perspectives. *Mol Med Rep*. 2019; 2019:20.
 217. Payne BAI, Chinnery PF. Mitochondrial dysfunction in aging: much progress but many unresolved questions. *Biochim Biophys Acta*. 2015; 2015:20.
 218. Kumsta C, Thamsen M, Jakob U. Effects of oxidative stress on behavior, physiology, and the redox thiol proteome of *Caenorhabditis elegans*. *Antioxid Redox Signal*. 2011; 2011:20.
 219. Hebbel RP, Osarogiagbon R, Kaul D. The endothelial biology of sickle cell disease: Inflammation and a chronic vasculopathy. *Microcirculation*. 2004; 2004:20.
 220. Kato GJ, Hebbel RP, Steinberg MH, Gladwin MT. Vasculopathy in sickle cell disease: Biology, pathophysiology, genetics, translational medicine, and new research directions. *Am J Hematol*. 2009; 2009:20.
 221. Singer ST, Ataga KI. Hypercoagulability in sickle cell disease and beta-thalassemia. *Curr Mol Med*. 2008; 2008:20.
 222. Sydow K, Munzel T. ADMA and oxidative stress. *Atheroscler Suppl*. 2003; 2003:20.
 223. Krajewski ML, Hsu LL, Gladwin MT. The proverbial chicken or the egg? Dissection of the role of cell-free hemoglobin versus reactive oxygen species in sickle cell pathophysiology. *Am J Physiol Heart Circ Physiol*. 2008; 295:H4-7.
 224. Morris CR, Suh JH, Hagar W, *et al.* Erythrocyte glutamine depletion, altered redox environment, and pulmonary hypertension in sickle cell disease. *Blood*. 2008; 2008:20.
 225. Nur E, Brandjes DP, Schnog JJ, *et al.* CURAMA Study Group. Plasma levels of advanced glycation end products are associated with haemolysis-related organ complications in sickle cell patients. *Br J Haematol*. 2010; 2010:20.
 226. Nur E, Biemond BJ, Otten HM, Brandjes DP, Schnog JJ; CURAMA Study Group. Oxidative stress in sickle cell disease; pathophysiology and potential implications for disease management. *Am J Hematol*. 2011; 2011:20.
 227. Atiku SM, Louise N, Kasozi DM. Severe oxidative stress in sickle cell disease patients with uncomplicated Plasmodium falciparum malaria in Kampala, Uganda. *BMC Infect Dis*. 2019; 2019:20.
 228. Biswal S, Rizwan H, Pal S, Sabnam S, Parida P, Pal A. Oxidative stress, antioxidant capacity, biomolecule damage, and inflammation symptoms of sickle cell disease in children. *Hematology*. 2019; 2019:20.
 229. Beetsch JW, Park TS, Dugan LL, Shah AR, Gidday JM. Xanthine oxidase-derived superoxide causes reoxygenation injury of ischemic cerebral endothelial cells. *Brain Res*. 1998; 1998:19.
 230. Solovey A, Gui L, Key NS, Hebbel RP. Tissue factor expression by endothelial cells in sickle cell anemia. *J Clin Invest*. 1998; 1998:19.
 231. Belcher JD, Mahaseth H, Welch TE, *et al.* Critical role of endothelial cell activation in hypoxia-induced vasoocclusion in transgenic sickle mice. *Am J Physiol Heart Circ Physiol*. 2005; 2005:20.
 232. Frenette PS. Sickle cell vaso-occlusion: multistep and multicellular paradigm. *Curr Opin Hematol*. 2002; 2002:20.
 233. Belhassen L, Pelle G, Sediame S, *et al.* Endothelial dysfunction in patients with sickle cell disease is related to selective impairment of shear stress-mediated vasodilation. *Blood*. 2001; 2001:20.
 234. Reiter CD, Wang X, Tanus-Santos JE, *et al.* Cell-free hemoglobin limits nitric oxide bioavailability in sickle-cell disease. *Nat Med*. 2002; 2002:20.
 235. Guzik TJ, Korb R, Mek-Guzik T. Nitric oxide and superoxide in inflammation and immune regulation. *J Physiol Pharmacol*. 2003; 2003:20.
 236. Villagra J, Shiva S, Hunter LA, Machado RF, Gladwin MT, Kato GJ. Platelet activation in patients with sickle disease, hemolysis-associated pulmonary hypertension, and nitric oxide scavenging by cell-free hemoglobin. *Blood*. 2007; 2007:20.
 237. Aslan M, Ryan TM, Adler B, *et al.* Oxygen radical inhibition of nitric oxide-dependent vascular function in sickle cell disease. *Proc Natl Acad Sci USA*. 2001; 2001:20.
 238. Banerjee T, Kuypers FA. Reactive oxygen species and phosphatidylserine externalization in murine sickle red cells. *Br J Haematol*. 2004; 2004:20.
 239. Cesquini M, Torsoni MA, Stoppa GR, Ogo SH. t-BOOH-induced oxidative damage in sickle red blood cells and the role of flavonoids. *Biomed Pharmacother*. 2003; 2003:20.
 240. Suzuki J, Umeda M, Sims PJ, Nagata S. Calcium-dependent phospholipid scrambling by TMEM16F. *Nature*. 2010; 2010:20.
 241. Neidinger NA, Larkin SK, Bhagat A, Victorino GP, Kuypers FA. Hydrolysis of phosphatidylserine-exposing red blood cells by secretory phospholipase A2 generates lysophosphatidic acid and results in vascular dysfunction. *J Biol Chem*. 2006; 2006:20.
 242. Thimmulappa RK, Chattopadhyay I, Rajasekaran S. Oxidative stress mechanisms in the pathogenesis of environmental lung diseases. In: Chakraborti S, Parinandi N, Ghosh R, Ganguly N, Chakraborti T, editors. *Oxidative Stress in Lung Diseases*. Singapore: Springer; 2020.
 243. Park HS, Kim SR, Lee YC. Impact of oxidative stress on lung diseases. *Respirology*. 2009; 2009:20.
 244. Kay AB. Asthma and inflammation. *J Allergy Clin Immunol*. 1991; 1991:19.

245. Bousquet J, Jeffery PK, Busse WW, Johnson M, Vignola AM. Asthma: From bronchoconstriction to airways inflammation and remodeling. *Am J Respir Crit Care Med.* 2000; 2000:20.
246. Russell RJ, Brightling C. Pathogenesis of asthma: implications for precision medicine. *Clin Sci.* 2017; 2017:20.
247. Henricks PA, Nijkamp FP. Reactive oxygen species as mediators in asthma. *Pulm Pharmacol Ther.* 2001; 2001:20.
248. Owen S, Pearson D, Suarez-Mendez V, O'Driscoll R, Woodcock A. Evidence of free-radical activity in asthma. *N Engl J Med.* 1991; 1991:19.
249. Adam L, Bouvier M, Jones TL. Nitric oxide modulates beta(2)-adrenergic receptor palmitoylation and signaling. *J Biol Chem.* 1999; 1999:19.
250. Calhoun WJ, Bush RK. Enhanced reactive oxygen species metabolism of airspace cells and airway inflammation following antigen challenge in human asthma. *J Allergy Clin Immunol.* 1990; 1990:19.
251. Calhoun WJ, Reed HE, Moest DR, Stevens CA. Enhanced superoxide production by alveolar macrophages and airspace cells, airway inflammation, and alveolar macrophage density changes follow segmental antigen bronchoprovocation in allergic subjects. *Am Rev Respir Dis.* 1992; 1992:19.
252. Schauer U, Leinhaas C, Jäger R, Rieger CH. Enhanced superoxide generation by eosinophils from asthmatic children. *Int Arch Allergy Immunol.* 1991; 1991:19.
253. Teramoto S, Shu CY, Ouchi Y, Fukuchi Y. Increased spontaneous production and generation of superoxide anion by blood neutrophils in patients with asthma. *J Asthma.* 1996; 1996:19.
254. Montuschi P, Collins JV, Ciabattoni G, *et al.* Exhaled 8-isoprostane as an *in vivo* biomarker of lung oxidative stress in patients with COPD and healthy smokers. *Am J Respir Crit Care Med.* 2000; 2000:20.
255. Hogg JC, Timens W. The pathology of chronic obstructive pulmonary disease. *Annu Rev Pathol.* 2009; 2009:20.
256. Hogg JC, Pare PD, Hackett TL. The contribution of small airway obstruction to the pathogenesis of chronic obstructive pulmonary disease. *Physiol Rev.* 2017; 2017:20.
257. Rangasamy T, Cho CY, Thimmulappa RK, *et al.* Genetic ablation of Nrf2 enhances susceptibility to cigarette smoke-induced emphysema in mice. *J Clin Invest.* 2004; 2004:20.
258. Singh A, Rangasamy T, Thimmulappa RK, *et al.* Glutathione peroxidase 2, the major cigarette smoke-inducible isoform of GPX in lungs, is regulated by Nrf2. *Am J Respir Cell Mol Biol.* 2006; 2006:20.
259. Yoshida T, Tudor RM. Pathobiology of cigarette smoke-induced chronic obstructive pulmonary disease. *Physiol Rev.* 2007; 2007:20.
260. Yoshida T, Mett I, Bhunia AK, *et al.* Rtp801, a suppressor of mTOR signaling, is an essential mediator of cigarette smoke-induced pulmonary injury and emphysema. *Nat Med.* 2010; 2010:20.
261. Thimmulappa RK, Gang X, Kim JH, Sussan TE, Witztum JL, Biswal S. Oxidized phospholipids impair pulmonary antibacterial defenses: evidence in mice exposed to cigarette smoke. *Biochem Biophys Res Commun.* 2012; 2012:20.
262. Tudor RM, Petrache I. Pathogenesis of chronic obstructive pulmonary disease. *J Clin Invest.* 2012; 2012:20.
263. Barnes PJ, Burney PG, Silverman EK, *et al.* Chronic obstructive pulmonary disease. *Nat Rev Dis Primers.* 2015; 2015:20.
264. Kirkham PA, Barnes PJ. Oxidative stress in COPD. *Chest.* 2013; 2013:20.
265. Dekhuijzen PN, Aben KK, Dekker I, *et al.* Increased exhalation of hydrogen peroxide in patients with stable and unstable chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 1996; 1996:19.
266. Maziak W, Loukides S, Culpitt S, Sullivan P, Kharitonov SA, Barnes PJ. Exhaled nitric oxide in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 1998; 1998:19.
267. Pratico D, Basili S, Vieri M, *et al.* Chronic obstructive pulmonary disease is associated with an increase in urinary levels of isoprostane F2alpha-III, an index of oxidant stress. *Am J Respir Crit Care Med.* 1998; 1998:19.
268. Rahman I, Morrison D, Donaldson K, MacNee W. Systemic oxidative stress in asthma, COPD and smokers. *Am J Respir Crit Care Med.* 1996; 1996:19.
269. Rahman I, Skwarska E, MacNee W. Attenuation of oxidant/antioxidant imbalance during treatment of exacerbations of chronic obstructive pulmonary disease. *Thorax.* 1997; 1997:19.
270. Fattman CL. Apoptosis in pulmonary fibrosis: too much or not enough? *Antioxid Redox Signal.* 2008; 2008:20.
271. Thannickal VJ, Fanburg BL. Activation of an H₂O₂-generating NADH oxidase in human lung fibroblasts by transforming growth factor beta 1. *J Biol Chem.* 1995; 1995:19.
272. Waghray M, Cui Z, Horowitz JC, *et al.* Hydrogen peroxide is a diffusible paracrine signal for the induction of epithelial cell death by activated myofibroblasts. *FASEB J.* 2005; 2005:20.
273. Sturrock A, Cahill B, Norman K, *et al.* Transforming growth factor-beta1 induces Nox4 NAD(P)H oxidase and reactive oxygen species dependent proliferation in human pulmonary artery smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol.* 2006; 2006:20.
274. Koli K, Myllarniemi M, Keski-Oja J, Kinnula VL. Transforming growth factor-β activation in the lung: focus on fibrosis and reactive oxygen species. *Antioxid Redox Signal.* 2008; 2008:20.
275. Behr J, Maier K, Krombach F, Adelman-Grill BC. Pathogenetic significance of reactive oxygen species in diffuse fibrosing alveolitis. *Am Rev Respir Dis.* 1991; 1991:19.
276. Lenz AG, Costabel U, Maier KL. Oxidized BAL fluid proteins in patients with interstitial lung diseases. *Eur Respir J.* 1996; 1996:19.
277. Rottoli P, Magi B, Cianti R, *et al.* Carbonylated proteins in bronchoalveolar lavage of patients with sarcoidosis, pulmonary fibrosis associated with systemic sclerosis and idiopathic pulmonary fibrosis. *Proteomics.* 2005; 2005:20.
278. Kang SK, Rabbani ZN, Folz RJ, *et al.* Overexpression of extracellular superoxide dismutase protects mice from radiation-induced lung injury. *Int J Radiat Oncol Biol Phys.* 2003; 2003:20.
279. Gao F, Kinnula VL, Myllarniemi M, Oury TD. Extracellular superoxide dismutase in pulmonary fibrosis. *Antioxid Redox Signal.* 2008; 2008:20.
280. Rahman I, Skwarska E, Henry M, *et al.* Systemic and pulmonary oxidative stress in idiopathic pulmonary fibrosis. *Free Radic Biol Med.* 1999; 1999:19.
281. Volm M, van Kaick G, Mattern J. Analysis of c-fos, c-erbB1, c-erbB2 and c-myc in primary lung carcinomas and their lymph node metastases. *Clin Exp Metastasis.* 1994; 1994:19.
282. Szabo E, Riffe ME, Steinberg SM, Birrer MJ, Linnoila RI. Altered cJUN expression: an early event in human lung carcinogenesis. *Cancer Res.* 1996; 1996:19.
283. Stewart ZA, Pietenpol JA. p53 signaling and cell cycle checkpoints. *Chem Res Toxicol.* 2001; 2001:20.

Cite this article: Ohiagu FO, Chikezie PC, Maduka TO, Chikezie CM, Nwaiwu O, Paudel KR. Antioxidants, Radical Scavenging, and Their Impact on Oxidative Stress. *Free Radicals and Antioxidants.* 2024;14(2):62-85.